

**Peconic Estuary Program 2004 Eelgrass (*Zostera marina*) Long-Term
Monitoring Program**

**Progress Report 6
September 21, 2005**

Submitted To:

**The Peconic Estuary Program Office
The Suffolk County Department of Health Services
Office of Ecology**

Submitted By:

**Christopher Pickerell
and
Stephen Schott**



Cornell University
Cooperative Extension
of Suffolk County
Marine Program

Summary

The Peconic Estuary Program's Long-Term Eelgrass Monitoring Program was continued by Cornell Cooperative Extension's Marine Program in 2004. The six monitoring beds were sampled during the period of 15 August 2004 to 22 August 2004. Divers conducted 60 quadrat counts of eelgrass shoot density and macroalgae percent cover at each monitoring site. Temperature data from data loggers and PEP Routine Marine Surface Water Monitoring Program were analyzed to elucidate differences in surface versus bottom temperatures and annual temperature trends. Significant changes in the general health and extent of the six monitoring sites were observed in 2004. Nine out of a total of 36 stations (6 stations per each of the 6 sites) no longer supported eelgrass within the 10 m of the station coordinates. Macroalgal percent cover has remained stable or declined in a majority of the monitoring sites. Areal extent has declined significantly in Bullhead Bay, Orient Harbor and Three Mile Harbor, with Southold Bay having experienced minor loss in area, but no noticeable retreat in its deep edge. The temperature data found little difference between surface and bottom temperature at the sites analyzed, but tracked annual and summer temperature trends well.

The significant decrease in eelgrass shoot densities at Bullhead Bay, Orient Harbor and Three Mile Harbor has been sudden and with little evidence of cause. Losses are not attributed to water quality or macroalgae competition, at this time, as these parameters have maintained healthy levels for several consecutive years. The lack of rhizome/root in the sediment at the sites suggest that physical disturbance was the probable factor in the loss, though the exact mechanism has not been identified. Possible mechanisms include, ice scour, anchor ice, changes in long-shore erosion/deposition patterns, and shellfishing activities. The temperature data was found to be a useful tool for use in monitoring annual trends and identifying localized periods of high water temperature, and it should be continued in subsequent monitoring efforts.

Eelgrass Introduction

The decline of eelgrass (*Zostera marina* L.) in the Peconic Estuary over the last 70 years has contributed to the degradation of the estuary as a whole. This submerged, marine plant is inextricably linked to the health of the Estuary, providing an important habitat in near shore waters for shellfish and finfish and a food source for organisms ranging from bacteria to waterfowl. To better manage this valuable resource, a baseline of data must be collected to identify trends in the health of the eelgrass meadows and plan for future work in the Peconic Estuary, such as conservation/management and restoration activities. The more data that is collected on the basic parameters of eelgrass, the better able the Peconic Estuary Program will be to implement policies to protect and nurture the resource.

The basic purpose of a monitoring program is to collect data on a regularly scheduled basis to develop a basic understanding of the ecology of the target species. Since its inception, the Peconic Estuary Program's Submerged Aquatic Vegetation Monitoring Program, contracted to Cornell Cooperative Extension's Marine Program, has focused on collecting data pertaining to the health of the eelgrass beds in the Peconic Estuary. The development of this program reflects an adaptation to the unique ecology and demography of the eelgrass in the Peconic estuary and varies significantly from other monitoring programs in the Chesapeake and other areas on the east coast, which tend to focus more on remote sensing techniques (i.e., aerial photography) for monitoring.

Methods

Table 1. The six reference eelgrass beds and the townships in which the beds are located.

| | |
|-------------------------|----------------|
| Bullhead Bay (BH) | Southampton |
| Gardiners Bay (GB) | Shelter Island |
| Northwest Harbor (NWH) | East Hampton |
| Orient Harbor (OH) | Southold |
| Southold Bay (SB) | Southold |
| Three Mile Harbor (TMH) | East Hampton |

The PEP SAV Monitoring Program includes six eelgrass beds located throughout the estuary and representing a range of environmental factors. The name and township location of each of the reference beds are listed in Table 1 and an aerial perspective of each site can be found in Appendix 1. Included with each image are the locations of the six sampling stations within the bed and the GPS coordinates for each station.

The monitoring program has evolved its methodologies from its inception in 1997, however, the basic parameter of eelgrass health, shoot density, has always been the focus of the program, thus allowing for comparisons between successive years. In the beginning, sampling consisted of the destructive collection of three (four in Bullhead Bay) 0.25 m² (50cm x 50cm) quadrats of eelgrass including below ground and above ground biomass that was returned to the laboratory for analysis. The sampling in 1998 and 1999 continued to utilize destructive sampling to collect data, however, sample size was increased to a total of twelve quadrats and there was a decrease in the size of the quadrats to 0.0625 m² (12.5 x 12.5 cm).

In 2000, the methodology for the

monitoring program was amended to increase the statistical significance of the data collected. The adjustments reflected an increase in the number of sampling stations per site (from 3 to 6), the number of replicate samples per station (from 4 to 10) and the size of the quadrats. However, the 2000 methodology included an increase number of destructively sampled quadrats (24 quadrats) for use in biomass estimations. The 2001 protocols maintained the higher number of replicate samples per bed (60 quadrats) but eliminated the destructive sampling aspect of the program. Beginning in 2004, water temperature was collected at several of the monitoring sites using submersible temperature loggers. The specific monitoring protocol for 2004 is outlined below.

Water Quality

Water quality data is supplied by the Suffolk County Department of Health Services. The data represents monthly to bimonthly sampling of various water quality parameters. The nitrogen-based data sets, nitrate/nitrite (NO_x), total nitrogen (TN), and total dissolved nitrogen (TDN), were analyzed for stations in or adjacent to the six, long-term monitoring sites and were incorporated into the long-term data set for trend analysis.

Water Temperature Monitoring

In an effort to better describe the relationship between water temperature and the life cycle of eelgrass, temperature loggers were deployed in several eelgrass beds in the Peconics, including 4 of the long-term monitoring sites. When possible, a surface logger (< 0.5 m from the surface) and a bottom logger (anchored to the bottom) were deployed at each site. The loggers were set to record temperature at

six-hour intervals.

The following sites have been chosen to pair with the existing PEP Routine Marine Surface Water Monitoring Program (RMSWMP) stations as well as existing eelgrass meadows: 109-Mill Creek (outer channel adjacent to eelgrass meadow), 126-Sag Harbor (outside breakwater adjacent to eelgrass meadow), 148-Bullhead Bay, and 118-Northwest Harbor (in existing eelgrass meadow).

Sites that are not directly associated with current PEP RMSWMP, but are essential based on existing monitoring and restoration efforts include Hallocks Bay (Orient), Hay Beach Point (Shelter Island), Sag Harbor Cove (directly behind Long Beach) and Orient Point (near Cross Island Ferry site).

The loggers, Onset Tidbit® and Onset StowAway®, were deployed in May 2004 and retrieved in September-October 2004, except for the logger in Sag Cove, which was left in place to provide a long-term temperature dataset at the eelgrass restoration area at this site.

The May-October deployment was designed to track the rise and fall of water temperature through 15°, a temperature thought to influence flowering and seed germination. This period also allows for peak water temperature, the most stressful time of the year for eelgrass, to be recorded.

Temperature data was exported from the loggers into spreadsheets. The data was analyzed and graphed using SigmaStat® and SigmaPlot® (SPSS Inc., 1997) software.

Eelgrass Monitoring

The monitor, for the 2004 season, was initiated on 15 August, 2004 and completed on 22 August, 2004.

Sampling at each site was distributed among six stations that have been referenced

using GPS. At each of the six stations, divers conducted a total of 10 random, replicate counts of eelgrass stem density and algal percent cover in 0.10 m² quadrats. Divers also made observations on blade lengths and overall health of plants that they observe. The divers stayed within a 10 meter radius of the GPS station point while conducting the survey. Algae within the quadrats were identified by genus and whether it was epiphytic or non-epiphytic on the eelgrass. Divers were careful not to disturb the eelgrass causing plants to be uprooted or otherwise damaged.

Data was incorporated into a spreadsheet and statistically analyzed using SigmaStat software (SPSS Inc., 1997). The trends, within sites, were analyzed by comparing the 2004 data with the data from the previous years.

Bed Delineation

For the 2004 season, the delineation for the deep edge was conducted using aerial photographs taken in Spring 2004. The aerial delineations were ground-truthed and found to accurately define the deep edge of most of the beds.

Genetics

Using microsatellite loci methodology to determine relatedness and diversity within and between these meadows we expect to be able to make some conclusions regarding the history of colonization and overall stability. An assessment of within bed diversity will allow us to determine the relative age and stability of these populations. This information will also allow us to better predict long-term survival of these meadows in the face of anthropogenic stressors. In addition to allowing us to better understand the history of extant meadows this information would

allow us to better determine sources of seeds and adult shoots for future restoration efforts.

The following sites have been chosen to serve as sample locations based on geographic distribution and relative hydrologic isolation: Shinnecock Bay (north shore east of canal), Bullhead Bay, Noyack Creek, Southold Bay, Sag Harbor, Hallocks Bay, Orient Point (west of ferry terminal), Long Island Sound (Mulford Point), Hog Creek, and Lake Montauk.

At each site, 25 samples will be taken at 6 successive 1 meter intervals along the major compass bearings (N,S,E & W) from a central point to determine within bed clonality/diversity. It is expected that some of these samples will yield identical genetic results, but until we test this hypothesis we cannot assume this to be the case.

Results

Statistical analysis reports are included in Appendices 5 and include basic descriptive statistics as well as one-way ANOVAs. *P*-values when, not stated, are included in Appendices 5-10, as well.

Water Quality

Water quality analysis is represented in Appendix 1. The graphs represent the mean annual concentrations of the three parameters measured (NO_x, TN, TDN) at or near the monitoring sites.

Bullhead Bay

Water quality continues to improve or remain stable in the Bullhead Bay system. Mean annual NO_x concentrations continue to decline in the system, while the TN and TDN have remained relatively stable (Appendix 1a.).

Gardiner's Bay

Gardiner's Bay has continued to maintain relatively low concentrations of nitrogen-based parameters. The site experienced a slight increase in NO_x concentration between 2002 and 2004, but analysis determined that it was statistically insignificant (Appendix 1b.). TN and TDN continue modest declines in concentration, though these changes were not significantly different between 2000 and 2004 (Appendix 1b.)

Northwest Harbor

All three water quality parameters continued to remain relatively stable in Northwest Harbor since 2000 (Appendix 1c.). While NO_x has remained consistent since 2001, TN and TDN have shown minor, and insignificant, decreases in concentration of both of these parameters (Appendix 1c.).

Orient Harbor

Orient harbor, as with the previous sites, has shown slight decreases in concentrations of NO_x, TN, and TDN since 2000 (Appendix 1d.). The only significant change in these parameters has been in TDN. The mean concentration has shown a significant ($p= 0.03$) decrease between 2000 and 2004 (Appendix 9).

Southold Bay

Appendix 1c represents the water quality data for Southold Bay. The data found no significant changes in parameter concentrations since 1999 at this site, though minor fluctuations in annual concentrations are evident (Appendix 1e).

Three Mile Harbor

The water quality in Three Mile Harbor has, like the other 5 sites, remained

relatively stable with regard to annual concentrations of NO_x, TN, and TDN (Appendix 1f). NO_x and TN, have not shown a significant change since 1999 and 2000, respectively. TDN, however, was found to have significantly decrease in concentration between 2000 and 2004 (Appendix 11).

Water Temperature Monitoring

The graphs for the water temperature data are included in Appendix 2. All temperature data for each site were included on one graph. In general, sites that included surface and bottom temperature loggers (Hallocks Bay and Southold), displayed little difference in temperatures, within the water column (Appendices 2band 2f). The PEP RMSWMP data shows a similar lack of significant temperature difference between surface and bottom samples in its dataset, as well.

The PEP RMSWMP data, when plotted with the temperature logger data, correlates well with the logger data (Appendices 2a,c,e,f). Several deviations between the logger data and the PEP RMSWMP data are evident (e.g. the high temperature recorded by PEP RMSWMP in August 2004 for Bullhead Bay), though may be an artifact of the software analysis.

Eelgrass Stem Density and Areal Extent

The basic descriptive statistics for the eelgrass stem densities for the 2004 season are represented in Table 2. Included in the table are the sample sizes (replicates), number of stations without eelgrass, mean stem density, and standard error of the

Table 2. Descriptive statistics for eelgrass stem density.

| Location | Sample Size (n) | # Stations w/ No Grass | Mean Stem Density (shoots/m ²) | Standard Error |
|-------------------------|-----------------|------------------------|--|----------------|
| Bullhead Bay (BH) | 60 | 4 | 126 | ±28.1 |
| Gardiner's Bay (GB) | 60 | 0 | 300 | ±26.4 |
| Northwest Harbor (NWH) | 60 | 0 | 291 | ±18.3 |
| Orient Harbor (OH) | 60 | 3 | 56 | ±14.6 |
| Southold Bay (SB) | 60 | 0 | 210 | ±23.3 |
| Three Mile Harbor (TMH) | 60 | 2 | 29 | ±6.1 |

means.

In the past, the stem density data was further analyzed by looking for differences in densities between beds. It was found in the 5-Year Trends Analysis Report (Pickerell and Schott, 2004), that trends analysis of within-bed variation was a more appropriate statistical measure than comparison between beds, and that analysis continues in this report. Appendix 3 includes trend analysis graphs of the shoot density data for the six monitoring sites.

Bullhead Bay

The Bullhead Bay eelgrass population suffered significant loss in areal coverage between 2002 and 2004, resulting in a significant decline in stem density in 2004. The mean stem density for 2004 was found to be 126 shoots/m² (Figure 2), representing a significant decrease ($p < 0.05$) in mean shoot density from 2002 (Appendix 3a). The areal extent of the bed declined to the western half of the bay and includes only 2 sampling stations (STA 3 and 4). The current extent of the bed is illustrated in Appendix 3a and represents a significant decrease in areal extent from 2002 to 2004.

Gardiners Bay

Gardiners Bay has continued to maintain a relatively stable eelgrass stem density. The 2004 mean stem density of 300 shoots/m² (Figure 2), while slightly lower than the 2002, does not represent a significant decrease (Appendix 3b). There has not been a significant change in eelgrass stem density in this bed since 2000 (Appendix 3b; Appendix 7), though there has been an overall significant decline in shoot density of approximately 200 shoots/m² from 1999 to 2004.

The areal extent of this bed continues to be dynamic from year to year. The “finger-like” projections representing the outermost extent of the bed are constantly shifting due to erosion and deposition. The aerial photograph in Appendix 3b, shows the patchy, irregular nature of the deep edge of this bed.

Northwest Harbor

Northwest Harbor has remained the most stable eelgrass bed in the program. There has not been a significant change in the eelgrass shoot density in this bed since the monitoring was initiated in 1997 (Appendix 3c and Appendix 8), though the data from 1997 to 1999 should be regarded carefully due to the small number of replicates. The

Table 3. Mean macroalgal percent coverage (m²).

| Eelgrass Bed | Percent Macroalgae Coverage |
|-------------------|-----------------------------|
| Bullhead Bay | < 1 |
| Gardiners Bay | 21.3 |
| Northwest Harbor | 81.4 |
| Orient Harbor | 1.1 |
| Southold Bay | 34.3 |
| Three Mile Harbor | 14.7 |

mean shoot density for 2004 (291 shoots/m²) did show a decline from 2002 (349 shoots/m²) (Appendix 3c), however the change was found to be insignificant after analysis (Appendix 8).

The areal extent of Northwest Harbor's eelgrass beds has shown moderate changes. However, it has been found that the deep edge delineated by Tiner (2003) was significantly underestimated when compared to the 2004 aerial photographs and groundtruthing (Appendix 4c).

Orient Harbor

Orient Harbor has seen the most drastic decline in its eelgrass stem density from previous years. In 2002, the mean stem density for Orient Harbor was 230 shoots/m² (Appendix 3d), but it decrease to only 56 shoots/m² (Figure 2) in 2004 (Appendix 3d). In 2004, 3 stations in this bed supported no eelgrass (STA 2,3 and 4), while 2 other stations (STA 1 and 6) showed reduced densities.

The areal extent of the bed is no longer distinct due to the heavy fragmentation occurring across the site, but the deep edge has noticeably receded from the prior depths of 7-9ft MLW (Appendix 4d).

Southold Bay

Southold Bay shoot density continued to show a decline from 1999 to 2004 (Appendix 3e). The decrease in shoot density from 2002 (384 shoots/m²) to 2004 (210 shoots/m²) was found to be a significant decline ($p < 0.05$) (Appendix 10).

The deep edge of the Southold Bay bed has remained relatively stable, though the areal extent of the bed has suffered some shrinkage along the eastern and western ends. (Appendix 4e).

Three Mile Harbor

Three Mile Harbor has seen a drastic decline in its eelgrass bed at the mouth of Hand's Creek. The mean shoot density declined from 135 shoots/m² in 2002, to just 29 shoots/m² in 2004 (Appendix 3f and Appendix 11). The 2004 survey found 2 stations devoid of grass, with 3 other stations supporting few plants.

The deep edge of this bed has not been affected by the above mentioned losses and remains at a depth of approximately 7-9ft MLW (Appendix 4f). However the areal extent of the bed has been significantly reduced with the loss of grass within two of the stations (Appendix 4f).

Algal Percent Cover

Algal percent cover was quantified for each quadrat within the six beds. Table 3 contains the mean percent coverage of macroalgae for each bed. Graphs for the individual sites are included in Appendix 5.

Bullhead Bay

Nonepiphytic, or drift, macroalgae has been a common component of the eelgrass community in Bullhead Bay since monitoring began in 1997. The percent cover of macroalgae at this site has been found to fluctuate from year to year, making a predictable trend difficult to identify

(Appendix 5a). The percent cover of macroalgae for 2004 was near zero (0.1%) (Table 3; Appendix 5a). With the loss of a large portion of the eelgrass bed, there has also been a decline in the macroalgae at the site. The dominate species encountered in Bullhead Bay continues to be the red, filamentous alga, *Spyridia filamentosa*, however, the areas once supporting eelgrass have shifted to bare bottom covered with films of diatoms and cyanobacteria.

Gardiner's Bay

Gardiner's Bay has supported a diverse, yet modest population of macroalgae. Since 2000, there has been no significant fluctuations in the percent cover of macroalgae at the site (Appendix 7) and the mean percent cover from 2002 to 2004 showed minimal change (Appendix 5b). The macroalgae population is greatly influenced by the fast currents that the site and the species that occur at the site can with the tides.

Northwest Harbor

Northwest Harbor has seen an increase in macroalgal percent cover since 2000 (Appendix 4c). With a mean percent cover of 81.4% (Table 3), the 2004 survey documented the highest percent algae cover, of any bed, that has been recorded since 2000, when analysis of the parameter was initiated (Appendix 5). This represents a substantial macroalgal population and, at this site, that population is dominated by *Spyridia filamentosa*, which almost forms a monoculture.

Orient Harbor

The macroalgae community in Orient Harbor has experienced a significant decline in 2004 (Appendix 5d). Though this site has not supported a large population of

macroalgae over the course of this monitoring program (Appendix 5d), the 1.1% cover (Table 3) recorded for 2004 is a significant decline from 2002 (Appendix 9).

Southold Bay

Percent cover of macroalgae did not show significant change from 2002 to 2004 (Appendix 5e; Appendix 10). Although macroalgae has continued to encroach upon the eastern and western edges of the bed, the overall percent cover has decreased since 2001 (Appendix 5e). The macroalgal species composition of this bed continues to be dominated by *Spyridia filamentosa*, though *Codium fragile* has become established in sections of the bed.

Three Mile Harbor

Three Mile Harbor has seen a significant decline in macroalgae between 2000 and 2004 (Appendix 11). As with Bullhead Bay and Orient Harbor, there may be a correlation with the loss of eelgrass affecting the macroalgae community. As with Bullhead Bay and Orient Harbor, the eelgrass loss has left bare muck or sand sediment that provides little attachment for macroalgae and typically supports only diatomaceous or cyanobacteria films.

Genetics

Given difficulties in identifying an appropriate lab to analyze the samples for the genetics study, no data was generated for this portion of the monitoring program. When the appropriate lab is identified, the data will be provided as an addendum to this report.

Discussion

Significant changes were observed in the 2004 monitoring season. While water

quality continued to improve or at least remain stable from 2002, several beds experienced significant declines in mean shoot density and losses in areal extent.

Long-Term Eelgrass Monitoring *Bullhead Bay*

Bullhead Bay exhibited a significant decline in eelgrass shoot density and areal extent. Although the shoot density had only decreased by 76 shoots/m² from 2002 (Appendix 3a), the site includes 4 stations that no longer supports eelgrass reducing the overall areal extent of the bed by 66-75%. The cause of the losses are not clear. It has been proposed that physical removal, via a disturbance mechanism (e.g. ice scour, erosion, dredging, etc.), results in the loss/removal of the entire plant (roots, rhizomes and shoots). However, if other factors were responsible for the decline of an eelgrass population (e.g. disease, temperature-induced mortality, anchor ice, etc.), then some of the plant material, primarily the rhizomes, would remain intact in the sediment for a limited number of years. Within Bullhead Bay, examination of the sediment at unvegetated stations revealed 2 stations that contained rhizomes in the sediment and 2 station that included no rhizomes in the sediment. This suggests that the loss of eelgrass at two of the stations (relatively shallow and compact sand sediment) may have experienced ice scour between 2002 and 2004, possible multiple times over two winters. The other two stations consist of soft, muck sediment and are of slightly greater depth. It is possible that these stations were spared the scouring, but encountered another stressor that resulted in mortality, but left behind the remnants of the plants.

It was also eluded to in the results that there may be a correlation between the

decline of the eelgrass and the subsequent decrease in macroalgae. The relationship between macroalgae and eelgrass includes the use of the eelgrass bed as an anchorage for drift macroalgae. In the specific case of Bullhead Bay, *Spyridia filamentosa*, being the dominant macroalgal species, grows intertwined in the eelgrass blades. Without this attachment, *Spyridia* has no anchorage and is subject to movement by the prevailing winds and currents.

Gardiner's Bay

Gardiner's Bay has supported a relatively stable eelgrass population with few changes over the years. The erosion/deposition caused by the fast currents at this site, continue to change the look and extent of the bed annually. However, it seems that natural propagation is able to keep pace with loss associated with erosion and other disturbances. The macroalgae community at this site is strongly influenced by the current and waves. A large proportion of the macroalgae encountered at the site are drift that became entangled in the eelgrass. When the current changes direction, much of the algae that accumulated on the previous tide, likely is dislodged and washed out of the bed, resulting in a high turn over rate of species and abundance.

One factor of concern at the Gardiner's Bay site is the continued boat traffic that travels over the bed to and from Greenport Harbor. Boats, large and small, tend to cut inside the green buoy marking the channel by as much as 200 yards. At MLW, there can be less than 3 ft of water over this bed and prop scars are common adjacent to the shoal on the Greenport side of the eelgrass bed. The addition of another navigation buoy closer to Hay Beach Point would likely reduce the traffic over this bed by directing boaters to the channel.

Northwest Harbor

There is little concern for the health and areal extent of the Northwest Harbor bed at this time. The eelgrass population seems to be stable and little change is evident in the bed.

Though there has been little change in the eelgrass population at this site, the macroalgal population, primarily *Spyridia filamentosa*, has continued to increase since 2000. Though this increase in a potential competitor is cause for some concern, there has been no observable impact on the eelgrass. It may be that because the eelgrass populations have remained so stable over the years, that the *Spyridia* has been able to take advantage of the relatively plentiful anchorage that the shoots provide and expand accordingly. Though the macroalgal population was relatively high at the time of monitoring, it is likely that it declines significantly after storms and during periods in which eelgrass shoots slough old leaves, reducing the overall population.

Orient Harbor

Orient Harbor has long been considered one of the most stable eelgrass beds in the Peconic Estuary. Based on aerial photography, the bed has changed little in 70 years, until 2004. The aerial photograph of the site, shows that the once stable beds has become patchy and the deep edge has retreated from its position of two years prior.

The cause of the losses in this bed are unclear. It had been suggested that bulkheading was to blame, however, the stations adjacent to the hardened shoreline do not show the loss in eelgrass that the stations adjacent to undeveloped shoreline display. Examination of the sediment at the unvegetated stations revealed no rhizomes, suggesting that the plants were completely

removed from the substrate by shellfishing or other physical disturbance. Ice scour may have been responsible for the loss in the shallow areas of the bed, but it is unlikely that it caused the loss in the deeper section of the bed, resulting in the shoreward retreat of the deep edge to only 5-6ft at MLW.

The decline macroalgae at this site is likely related to the loss of an anchoring substrate, in this case, eelgrass. The mechanism for this was detailed above and is the most likely causative factor in the decrease in algal percent cover as nitrogen concentrations have not significantly decreased at this site.

Southold Bay

The shoot density at Southold Bay continues to decline, though there is not a clear cause for this trend. Although this site was originally chosen due to the opinion that it was a “bed in decline,” it has proven to be more stable, to date, than other beds in the program. Water quality has improved at the site, likely relieving some stress on the eelgrass population, though water clarity may still be a stressor, as the site includes two, high-traffic boat channels that result in considerable turbidity (Pickerell and Schott, personal observations).

Changes in macroalgal abundance (percent cover) have not shown significant change since 2002, though encroachment by *Codium fragile* in areas that once supported eelgrass has increased and may prevent the bed from recolonizing lost areas. The current, relatively low abundance of macroalgae in this bed should provide more favorable growing conditions for the eelgrass.

Three Mile Harbor

The eelgrass bed in Three Mile Harbor is in rapid decline. Though it has only lost

eelgrass in two stations, the shoot densities for the other stations have shown a marked decrease. The eelgrass growing near the deep edge are rooted in loose muck in 7-9 ft of water (MLW) and are easily uprooted by any disturbance that could fluidize the sediment. Turbidity is commonly high at these stations resulting in decreased light for growth and low shoot densities. In the shallower stations, the sediments are primarily sand, resulting in less turbidity and better anchorage. However, these stations experience more anthropogenic disturbances, such as mooring chain scour and prop scarring due to the mooring field and waterskiing area that are included in the confines of the bed. These continued activities in the eelgrass bed could lead to significant decline in the areal extent and overall health of the bed.

Macroalgae has not be a major concern for this bed. The percent cover has declined since 2000 and is significantly lower than other eelgrass beds in the program. It is likely that, as eelgrass densities continue to decline in Three Mile Harbor, the macroalgae will also decline due to loss of substrate for anchorage/attachment.

Overview

Significant changes in the general health and extent of the six monitoring sites were observed in 2004. Nine out of a total of 36 stations (6 stations per each of the 6 sites) no longer supported eelgrass within the 10 m of the station coordinates. That is up from only 2 stations in 2002. The three sites that represent these losses were Bullhead Bay (4 unvegetated stations), Orient Harbor (3 unvegetated stations) and Three Mile Harbor (2 unvegetated stations). These three beds represent an average loss of 118 shoots/m². Southold Bay also showed a significant decline in shoot density from

2002, with an approximate loss of 174 shoots/m².(Gardiner's Bay and Northwest Harbor continue to maintain stable shoot densities.

Areal extent has declined significantly in Bullhead Bay, Orient Harbor and Three Mile Harbor, and Southold Bay has experienced minor loss in area, but no noticeable retreat in its deep edge. Gardiner's Bay continues to be dynamic in its areal extent and deep edge, making it difficult to assess actual loss/gain at the site. Northwest Harbor is relatively unchanged from 2002 in extent and deep edge.

Macroalgal percent cover has remained stable or declined in a majority of the monitoring sites. The exception to this was Northwest Harbor, which saw an almost 20% increase in mean percent cover from 2002. Even with this increase in macroalgae, there was no significant decline in the eelgrass shoot density at this site.

The mechanism(s) of the reported losses have not been identified, but it has been suggested that the losses were the result of physical disturbances, rather than a biological agent or physiological event, due to the lack or rhizome/root remains in the sediment at the sites. Events that could explain these losses include ice scour, anchor ice, changes in local long-shore erosion/deposition patterns and dredging/shellfishing activities. The Peconic Estuary has experienced two cold winters since 2002 that resulted in most of the Estuary experiencing some level of freezing. This is of special concern in shallow areas, like eelgrass meadows, where the ice layers could sit on the bottom at low tides and crush the eelgrass or scour the sediment as it is moved by winds and tides. This is the likely cause of the loss of eelgrass in stations 5 and 6 in Bullhead Bay. Their location, in the south-southeast section

of the bay make them susceptible to ice being piled up in the shallow waters by prevailing winter winds. Anchor ice, though a recognized event in estuarine and riverine waters, has not been the focus of much research and therefore it is unclear of the extent of damage this event could cause to seagrasses. Changes in long-shore currents, though not a probable cause for loss, still holds the potential to damage an eelgrass bed by changing the erosion/deposition dynamics of the site. These currents may be influenced by changes in shoreline structure (i.e., hardening shorelines) and could require several years for the change to significantly manifest itself. Lastly are dredging and shellfishing. Both are human-based sources of disturbance and may influence any or all of the monitoring beds at some point in time. Although anecdotal evidence has suggested that shellfishing impacts to meadows in the Peconic Estuary (e.g. Hallocks Bay), it is unlikely, due to the area of the eelgrass loss at the three sites, that the damage observed was caused by shellfishing activities, and no evidence of dredging was observed at these monitoring sites. The situation requires continued monitoring and the 2005 monitoring season may shed light on the losses observed in 2004.

Water Temperature Monitoring

The analysis of water temperature data from deployed temperature loggers and from the PEP RMSWMP has resulted in several conclusions. The PEP RMSWMP, though limited in the number of samples, describes a relatively accurate annual trend in water temperature for the Peconic Estuary, when compared to the continual, 4 -month dataset. While the PEP RMSWMP data is adequate for determining large time-scale (i.e., monthly) trends in temperature, it does not

have the precision to illustrate the temperature dynamics on a smaller time-scale, like the logger data.

The analysis of surface versus bottom temperature data found no overall, significant differences between the surface and bottom temperatures at the indicated sites. The lack of significant differences between surface and bottom temperatures is likely due to the shallow nature of the sites. The depth at these sites, less than 3 meters, does not represent a significant depth for a thermocline to be established, or if a thermocline does occur, to be easily disrupted and mixed. While deploying loggers at sites with a greater depth may result in a significant difference between surface and bottom temperatures, these sites would not necessarily reflect the conditions experienced in the Estuary's eelgrass beds, which are relatively shallow.


The benefits of these 2 datasets include the prediction/planning of restoration activities (i.e., timing of seed collection and germination). The data could also be used to explain acute/chronic changes, especially eelgrass loss in the Estuary, by identifying localized periods of high water temperature and/or long-term shifts in temperatures due to climatic events (e.g. global warming).

References


Pickerell, C. and S. Schott, 2004. Peconic Estuary Program Long Term Eelgrass Monitoring Program - Eelgrass Trends Analysis Report: 1997-2002. Prepared for the Peconic Estuary Program, SCDHS, Department of Ecology.

Appendix 1. Graphs plotting the 2004 annual mean NO_x, TN, and TDN, based on the SCDHS Water Quality Data for the six long-term eelgrass monitoring sites. (Concentrations represent annual means in mg L⁻¹).


a) Bullhead Bay




b) Gardiner's Bay




c) Northwest Harbor




d) Orient Harbor




e) Southold Bay





f) Three Mile Harbor





Appendix 2. Water temperature graphs for selected sites within the Peconic Estuary. Datasets include continuous logging by temperature loggers deployed by CCE and data from PEP RMSWMP.
















Appendix 3. Graphs of the mean eelgrass shoot densities for the six long-term monitoring sites. (Shoot density is expressed as shoots m^{-2}).


a) Bullhead Bay




b) Gardiner's Bay




c) Northwest Harbor




d) Orient Harbor



e) Southold Harbor




f) Three Mile Harbor




Appendix 4. Aerial photographs, with deep edge delineations, of the six monitoring sites for 2004. Monitoring stations are indicated by numbers (1-6) for each site.


a) Bullhead Bay



b) Gardiner's Bay




c) Northwest Harbor



d) Orient Harbor



e) Southold Bay




f) Three Mile Harbor




Appendix 5. Graphs representing the mean Percent macroalgal cover at the six sites from 2000 to 2004.


a) Bullhead Bay




b) Gardiner's Bay




c) Northwest Harbor




d) Orient Harbor



e) Southold Bay



f) Three Mile Harbor



Appendix 6. The statistical reports generated for all of the analyses conducted in the synthesis of 2004 Long-Term Monitoring Report for Bullhead Bay.

Descriptive Statistics:

Data source: BB Water Quality Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|-----------|------|---------|--------|---------|------------|--------------|
| BB 98 NOx | 16 | 0 | 0.0051 | 0.00025 | 0.000063 | 0.00013 |
| BB 99 NOX | 17 | 0 | 0.0057 | 0.0022 | 0.00053 | 0.0011 |
| BB 00 NOx | 20 | 0 | 0.011 | 0.011 | 0.0025 | 0.0051 |
| BB 01 NOx | 22 | 0 | 0.037 | 0.032 | 0.0067 | 0.014 |
| BB 02 NOx | 26 | 0 | 0.027 | 0.030 | 0.0058 | 0.012 |
| BB 04 NOx | 12 | 0 | 0.013 | 0.015 | 0.0042 | 0.0093 |
| BB 00 TN | 10 | 0 | 0.30 | 0.061 | 0.019 | 0.044 |
| BB 01 TN | 22 | 0 | 0.23 | 0.078 | 0.017 | 0.035 |
| BB 02 TN | 26 | 0 | 0.21 | 0.074 | 0.015 | 0.030 |
| BB 04 TN | 12 | 0 | 0.21 | 0.20 | 0.058 | 0.13 |
| BB 00 TDN | 9 | 0 | 0.28 | 0.072 | 0.024 | 0.056 |
| BB 01 TDN | 22 | 0 | 0.21 | 0.082 | 0.018 | 0.037 |
| BB 02 TDN | 26 | 0 | 0.18 | 0.068 | 0.013 | 0.028 |
| BB 04 TDN | 12 | 0 | 0.18 | 0.20 | 0.057 | 0.12 |

| Column | Range | Max | Min | Median | 25% | 75% |
|-----------|---------|--------|--------|--------|--------|--------|
| BB 98 NOx | 0.00100 | 0.0060 | 0.0050 | 0.0050 | 0.0050 | 0.0050 |
| BB 99 NOX | 0.0090 | 0.014 | 0.0050 | 0.0050 | 0.0050 | 0.0052 |
| BB 00 NOx | 0.037 | 0.042 | 0.0050 | 0.0055 | 0.0050 | 0.012 |
| BB 01 NOx | 0.10 | 0.11 | 0.0050 | 0.028 | 0.0090 | 0.056 |
| BB 02 NOx | 0.11 | 0.12 | 0.0050 | 0.014 | 0.0070 | 0.040 |
| BB 04 NOx | 0.053 | 0.058 | 0.0050 | 0.0095 | 0.0050 | 0.014 |
| BB 00 TN | 0.17 | 0.39 | 0.22 | 0.30 | 0.27 | 0.35 |
| BB 01 TN | 0.25 | 0.38 | 0.13 | 0.20 | 0.19 | 0.27 |
| BB 02 TN | 0.29 | 0.37 | 0.080 | 0.19 | 0.16 | 0.22 |
| BB 04 TN | 0.64 | 0.69 | 0.050 | 0.13 | 0.060 | 0.34 |
| BB 00 TDN | 0.18 | 0.38 | 0.20 | 0.29 | 0.21 | 0.35 |
| BB 01 TDN | 0.27 | 0.39 | 0.12 | 0.18 | 0.16 | 0.24 |
| BB 02 TDN | 0.24 | 0.33 | 0.090 | 0.17 | 0.12 | 0.20 |
| BB 04 TDN | 0.56 | 0.61 | 0.050 | 0.080 | 0.050 | 0.29 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|-----------|----------|----------|-----------|-----------|-------|----------------|
| BB 98 NOx | 4.00 | 16.00 | 0.54 | <0.001 | 0.081 | 0.00041 |
| BB 99 NOX | 3.91 | 15.70 | 0.39 | <0.001 | 0.097 | 0.00063 |
| BB 00 NOx | 2.14 | 3.79 | 0.31 | <0.001 | 0.22 | 0.0048 |
| BB 01 NOx | 0.88 | -0.10 | 0.18 | 0.052 | 0.82 | 0.051 |
| BB 02 NOx | 1.78 | 2.64 | 0.27 | <0.001 | 0.70 | 0.041 |
| BB 04 NOx | 3.07 | 10.01 | 0.37 | <0.001 | 0.16 | 0.0044 |
| BB 00 TN | 0.026 | -1.26 | 0.15 | 0.676 | 3.03 | 0.95 |
| BB 01 TN | 0.97 | -0.13 | 0.23 | 0.004 | 5.00 | 1.26 |
| BB 02 TN | 0.93 | 0.43 | 0.20 | 0.007 | 5.33 | 1.23 |
| BB 04 TN | 1.48 | 1.75 | 0.23 | 0.091 | 2.52 | 0.97 |
| BB 00 TDN | 0.23 | -1.68 | 0.18 | 0.503 | 2.55 | 0.76 |
| BB 01 TDN | 1.10 | -0.0034 | 0.24 | 0.002 | 4.64 | 1.12 |
| BB 02 TDN | 0.90 | 0.018 | 0.15 | 0.112 | 4.68 | 0.96 |

BB 04 TDN 1.41 0.62 0.28 0.012 2.19 0.82

One Way Analysis of Variance

Data source: BB Water Quality Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: BB in Water Quality Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|-----------|----|---------|--------|--------|--------|
| BB 98 NOx | 16 | 0 | 0.0050 | 0.0050 | 0.0050 |
| BB 99 NOx | 17 | 0 | 0.0050 | 0.0050 | 0.0052 |
| BB 00 NOx | 20 | 0 | 0.0055 | 0.0050 | 0.012 |
| BB 01 NOx | 22 | 0 | 0.028 | 0.0090 | 0.056 |
| BB 02 NOx | 26 | 0 | 0.014 | 0.0070 | 0.040 |
| BB 04 NOx | 12 | 0 | 0.0095 | 0.0050 | 0.014 |

H = 43.24 with 5 degrees of freedom. (P = <0.001)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = <0.001)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|------------------------|---------------|------|--------|
| BB 01 NOx vs BB 98 NOx | 53.11 | 4.93 | Yes |
| BB 01 NOx vs BB 99 NOx | 46.86 | 4.43 | Yes |
| BB 01 NOx vs BB 00 NOx | 30.64 | 3.03 | Yes |
| BB 01 NOx vs BB 04 NOx | 21.93 | 1.86 | No |
| BB 01 NOx vs BB 02 NOx | 8.98 | 0.95 | No |
| BB 02 NOx vs BB 98 NOx | 44.12 | 4.24 | Yes |
| BB 02 NOx vs BB 99 NOx | 37.87 | 3.71 | Yes |
| BB 02 NOx vs BB 00 NOx | 21.65 | 2.22 | No |
| BB 02 NOx vs BB 04 NOx | 12.95 | 1.13 | No |
| BB 04 NOx vs BB 98 NOx | 31.18 | 2.49 | No |
| BB 04 NOx vs BB 99 NOx | 24.93 | 2.02 | No |
| BB 04 NOx vs BB 00 NOx | 8.71 | 0.73 | No |
| BB 00 NOx vs BB 98 NOx | 22.47 | 2.04 | No |
| BB 00 NOx vs BB 99 NOx | 16.22 | 1.50 | No |
| BB 99 NOx vs BB 98 NOx | 6.25 | 0.55 | No |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

One Way Analysis of Variance

Data source: BB Water Quality Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: BB in Water Quality Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|----------|----|---------|--------|-------|------|
| BB 00 TN | 10 | 0 | 0.30 | 0.27 | 0.35 |
| BB 01 TN | 22 | 0 | 0.20 | 0.19 | 0.27 |
| BB 02 TN | 26 | 0 | 0.19 | 0.16 | 0.22 |
| BB 04 TN | 12 | 0 | 0.13 | 0.060 | 0.34 |

H = 12.56 with 3 degrees of freedom. (P = 0.006)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = 0.006)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|----------------------|---------------|------|--------|
| BB 00 TN vs BB 04 TN | 28.17 | 3.23 | Yes |
| BB 00 TN vs BB 02 TN | 23.42 | 3.09 | Yes |
| BB 00 TN vs BB 01 TN | 17.41 | 2.24 | No |
| BB 01 TN vs BB 04 TN | 10.76 | 1.47 | No |
| BB 01 TN vs BB 02 TN | 6.01 | 1.02 | No |
| BB 02 TN vs BB 04 TN | 4.74 | 0.67 | No |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

One Way Analysis of Variance

Data source: BB Water Quality Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: BB in Water Quality Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|-----------|----|---------|--------|-------|------|
| BB 00 TDN | 9 | 0 | 0.29 | 0.21 | 0.35 |
| BB 01 TDN | 22 | 0 | 0.18 | 0.16 | 0.24 |
| BB 02 TDN | 26 | 0 | 0.17 | 0.12 | 0.20 |
| BB 04 TDN | 12 | 0 | 0.080 | 0.050 | 0.29 |

H = 13.69 with 3 degrees of freedom. (P = 0.003)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = 0.003)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|------------------------|---------------|------|--------|
| BB 00 TDN vs BB 04 TDN | 30.68 | 3.47 | Yes |
| BB 00 TDN vs BB 02 TDN | 22.94 | 2.96 | Yes |
| BB 00 TDN vs BB 01 TDN | 15.92 | 2.01 | No |
| BB 01 TDN vs BB 04 TDN | 14.76 | 2.05 | No |
| BB 01 TDN vs BB 02 TDN | 7.02 | 1.21 | No |
| BB 02 TDN vs BB 04 TDN | 7.74 | 1.11 | No |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

Descriptive Statistics:

Data source: BB in Eelgrass Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|---------|------|---------|--------|---------|------------|--------------|
| BB 1997 | 4 | 0 | 710.00 | 392.32 | 196.16 | 624.27 |
| BB 1998 | 12 | 0 | 620.00 | 387.15 | 111.76 | 245.98 |
| BB 1999 | 12 | 0 | 548.00 | 271.97 | 78.51 | 172.80 |
| BB 2000 | 60 | 0 | 301.17 | 200.09 | 25.83 | 51.69 |
| BB 2001 | 60 | 0 | 150.17 | 138.66 | 17.90 | 35.82 |
| BB 2002 | 60 | 0 | 201.17 | 109.19 | 14.10 | 28.21 |
| BB 2004 | 60 | 0 | 125.50 | 217.85 | 28.12 | 56.28 |

| Column | Range | Max | Min | Median | 25% | 75% |
|---------|---------|---------|--------|--------|--------|--------|
| BB 1997 | 920.00 | 1264.00 | 344.00 | 616.00 | 460.00 | 960.00 |
| BB 1998 | 1184.00 | 1296.00 | 112.00 | 424.00 | 368.00 | 976.00 |
| BB 1999 | 944.00 | 1136.00 | 192.00 | 496.00 | 368.00 | 672.00 |
| BB 2000 | 880.00 | 930.00 | 50.00 | 250.00 | 155.00 | 385.00 |
| BB 2001 | 820.00 | 820.00 | 0.00 | 130.00 | 55.00 | 210.00 |
| BB 2002 | 450.00 | 450.00 | 0.00 | 180.00 | 120.00 | 270.00 |
| BB 2004 | 870.00 | 870.00 | 0.00 | 0.00 | 0.00 | 180.00 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|---------|----------|----------|-----------|-----------|----------|----------------|
| BB 1997 | 1.31 | 2.33 | 0.30 | 0.209 | 2840.00 | 2478144.00 |
| BB 1998 | 0.42 | -1.27 | 0.27 | 0.016 | 7440.00 | 6261504.00 |
| BB 1999 | 0.92 | 0.69 | 0.21 | 0.155 | 6576.00 | 4417280.00 |
| BB 2000 | 1.34 | 1.50 | 0.15 | 0.002 | 18070.00 | 7804300.00 |
| BB 2001 | 2.21 | 8.59 | 0.14 | 0.005 | 9010.00 | 2487300.00 |
| BB 2002 | 0.18 | -0.63 | 0.094 | 0.210 | 12070.00 | 3131500.00 |
| BB 2004 | 1.73 | 2.06 | 0.38 | <0.001 | 7530.00 | 3745100.00 |

One Way Analysis of Variance

Data source: BB in Eelgrass Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: BB in Eelgrass Trend analysis

| Group | N | Missing | Median | 25% | 75% |
|---------|----|---------|--------|--------|--------|
| BB 1997 | 4 | 0 | 616.00 | 460.00 | 960.00 |
| BB 1998 | 12 | 0 | 424.00 | 368.00 | 976.00 |
| BB 1999 | 12 | 0 | 496.00 | 368.00 | 672.00 |
| BB 2000 | 60 | 0 | 250.00 | 155.00 | 385.00 |
| BB 2001 | 60 | 0 | 130.00 | 55.00 | 210.00 |
| BB 2002 | 60 | 0 | 180.00 | 120.00 | 270.00 |
| BB 2004 | 60 | 0 | 0.00 | 0.00 | 180.00 |

H = 87.37 with 6 degrees of freedom. (P = <0.001)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = <0.001)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|--------------------|---------------|------|--------|
| BB 1997 vs BB 2004 | 162.41 | 4.06 | Yes |
| BB 1997 vs BB 2001 | 135.47 | 3.38 | Yes |
| BB 1997 vs BB 2002 | 104.66 | 2.61 | No |
| BB 1997 vs BB 2000 | 76.12 | 1.90 | No |
| BB 1997 vs BB 1998 | 21.67 | 0.48 | No |
| BB 1997 vs BB 1999 | 16.63 | 0.37 | No |
| BB 1999 vs BB 2004 | 145.78 | 5.95 | Yes |
| BB 1999 vs BB 2001 | 118.84 | 4.85 | Yes |
| BB 1999 vs BB 2002 | 88.03 | 3.59 | No |
| BB 1999 vs BB 2000 | 59.49 | 2.43 | No |
| BB 1999 vs BB 1998 | 5.04 | 0.16 | No |
| BB 1998 vs BB 2004 | 140.74 | 5.74 | Yes |
| BB 1998 vs BB 2001 | 113.80 | 4.64 | Yes |
| BB 1998 vs BB 2002 | 82.99 | 3.39 | No |
| BB 1998 vs BB 2000 | 54.45 | 2.22 | No |
| BB 2000 vs BB 2004 | 86.29 | 6.10 | Yes |
| BB 2000 vs BB 2001 | 59.35 | 4.19 | Yes |
| BB 2000 vs BB 2002 | 28.54 | 2.02 | No |
| BB 2002 vs BB 2004 | 57.75 | 4.08 | Yes |
| BB 2002 vs BB 2001 | 30.81 | 2.18 | No |
| BB 2001 vs BB 2004 | 26.94 | 1.90 | No |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

Descriptive Statistics:

Data source: BB Algae Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|--------|------|---------|------|---------|------------|--------------|
|--------|------|---------|------|---------|------------|--------------|

| | | | | | | |
|---------|----|---|-------|-------|-------|-------|
| BB 2000 | 24 | 0 | 18.96 | 20.11 | 4.10 | 8.49 |
| BB 2001 | 60 | 0 | 23.27 | 33.11 | 4.28 | 8.55 |
| BB 2002 | 60 | 0 | 56.42 | 38.98 | 5.03 | 10.07 |
| BB 2004 | 60 | 0 | 0.100 | 0.66 | 0.085 | 0.17 |

| Column | Range | Max | Min | Median | 25% | 75% |
|---------|--------|--------|------|--------|-------|--------|
| BB 2000 | 75.00 | 75.00 | 0.00 | 15.00 | 2.50 | 25.00 |
| BB 2001 | 100.00 | 100.00 | 0.00 | 5.00 | 0.00 | 50.00 |
| BB 2002 | 100.00 | 100.00 | 0.00 | 50.00 | 10.00 | 100.00 |
| BB 2004 | 5.00 | 5.00 | 0.00 | 0.00 | 0.00 | 0.00 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|---------|----------|----------|-----------|-----------|---------|----------------|
| BB 2000 | 1.50 | 2.15 | 0.26 | <0.001 | 455.00 | 17925.00 |
| BB 2001 | 1.25 | 0.14 | 0.29 | <0.001 | 1396.00 | 97176.00 |
| BB 2002 | -0.14 | -1.60 | 0.20 | <0.001 | 3385.00 | 280625.00 |
| BB 2004 | 7.34 | 55.27 | 0.53 | <0.001 | 6.00 | 26.00 |

One Way Analysis of Variance

Data source: BB Algae Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: BB in Algae Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|---------|----|---------|--------|-------|--------|
| BB 2000 | 24 | 0 | 15.00 | 2.50 | 25.00 |
| BB 2001 | 60 | 0 | 5.00 | 0.00 | 50.00 |
| BB 2002 | 60 | 0 | 50.00 | 10.00 | 100.00 |
| BB 2004 | 60 | 0 | 0.00 | 0.00 | 0.00 |

H = 98.12 with 3 degrees of freedom. (P = <0.001)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = <0.001)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|--------------------|---------------|------|--------|
| BB 2002 vs BB 2004 | 100.09 | 9.29 | Yes |
| BB 2002 vs BB 2001 | 50.29 | 4.67 | Yes |
| BB 2002 vs BB 2000 | 39.20 | 2.75 | Yes |
| BB 2000 vs BB 2004 | 60.90 | 4.27 | Yes |
| BB 2000 vs BB 2001 | 11.10 | 0.78 | No |
| BB 2001 vs BB 2004 | 49.80 | 4.62 | Yes |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

Appendix 7. The statistical reports generated for all of the analyses conducted in the synthesis of 2004 Long-Term Monitoring Report for Gardiner's Bay.

Descriptive Statistics:

Data source: GB Water Quality Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|-----------------------|------|---------|--------|----------------|----------------|----------------|
| GB 97 NO _x | 16 | 0 | 0.0054 | 0.0015 | 0.00037 | 0.00080 |
| GB 98 NO _x | 15 | 0 | 0.0050 | 0.000000000088 | 0.000000000023 | 0.000000000049 |
| GB 99 NO _x | 13 | 0 | 0.0072 | 0.0056 | 0.0016 | 0.0034 |
| GB 00 NO _x | 18 | 0 | 0.022 | 0.022 | 0.0051 | 0.011 |
| GB 01 NO _x | 7 | 0 | 0.038 | 0.022 | 0.0082 | 0.020 |
| GB 02 NO _x | 16 | 0 | 0.013 | 0.011 | 0.0029 | 0.0061 |
| GB 04 NO _x | 8 | 0 | 0.038 | 0.035 | 0.012 | 0.029 |
| GB 00 TN | 12 | 0 | 0.26 | 0.081 | 0.023 | 0.051 |
| GB 01 TN | 7 | 0 | 0.25 | 0.097 | 0.037 | 0.090 |
| GB 02 TN | 16 | 0 | 0.20 | 0.084 | 0.021 | 0.045 |
| GB 04 TN | 8 | 0 | 0.19 | 0.15 | 0.051 | 0.12 |
| GB 00 TDN | 12 | 0 | 0.24 | 0.084 | 0.024 | 0.053 |
| GB 01 TDN | 7 | 0 | 0.23 | 0.10 | 0.039 | 0.094 |
| GB 02 TDN | 16 | 0 | 0.19 | 0.091 | 0.023 | 0.048 |
| GB 04 TDN | 8 | 0 | 0.13 | 0.12 | 0.041 | 0.097 |

| Column | Range | Max | Min | Median | 25% | 75% |
|-----------------------|--------|--------|--------|--------|--------|--------|
| GB 97 NO _x | 0.0060 | 0.011 | 0.0050 | 0.0050 | 0.0050 | 0.0050 |
| GB 98 NO _x | 0.00 | 0.0050 | 0.0050 | 0.0050 | 0.0050 | 0.0050 |
| GB 99 NO _x | 0.018 | 0.023 | 0.0050 | 0.0050 | 0.0050 | 0.0050 |
| GB 00 NO _x | 0.054 | 0.059 | 0.0050 | 0.0090 | 0.0050 | 0.045 |
| GB 01 NO _x | 0.053 | 0.059 | 0.0060 | 0.045 | 0.017 | 0.057 |
| GB 02 NO _x | 0.033 | 0.038 | 0.0050 | 0.0060 | 0.0050 | 0.018 |
| GB 04 NO _x | 0.10 | 0.11 | 0.0050 | 0.028 | 0.011 | 0.055 |
| GB 00 TN | 0.24 | 0.36 | 0.12 | 0.27 | 0.19 | 0.34 |
| GB 01 TN | 0.24 | 0.36 | 0.12 | 0.20 | 0.19 | 0.35 |
| GB 02 TN | 0.31 | 0.39 | 0.080 | 0.16 | 0.14 | 0.27 |
| GB 04 TN | 0.39 | 0.44 | 0.050 | 0.14 | 0.075 | 0.31 |
| GB 00 TDN | 0.24 | 0.35 | 0.11 | 0.25 | 0.16 | 0.32 |
| GB 01 TDN | 0.24 | 0.35 | 0.11 | 0.19 | 0.16 | 0.34 |
| GB 02 TDN | 0.30 | 0.38 | 0.080 | 0.15 | 0.13 | 0.28 |
| GB 04 TDN | 0.32 | 0.37 | 0.050 | 0.075 | 0.050 | 0.20 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|-----------------------|----------|----------|-----------|-----------|-------|----------------|
| GB 97 NO _x | 4.00 | 16.00 | 0.54 | <0.001 | 0.086 | 0.00050 |
| GB 98 NO _x | 0.00 | -2.33 | 0.50 | <0.001 | 0.075 | 0.00038 |
| GB 99 NO _x | 2.47 | 5.41 | 0.50 | <0.001 | 0.094 | 0.0011 |
| GB 00 NO _x | 0.87 | -1.11 | 0.27 | 0.001 | 0.39 | 0.016 |
| GB 01 NO _x | -0.70 | -1.36 | 0.20 | 0.480 | 0.27 | 0.013 |
| GB 02 NO _x | 1.42 | 0.65 | 0.31 | <0.001 | 0.20 | 0.0045 |
| GB 04 NO _x | 1.27 | 1.43 | 0.19 | 0.491 | 0.30 | 0.020 |
| GB 00 TN | -0.21 | -1.37 | 0.17 | 0.371 | 3.08 | 0.86 |
| GB 01 TN | 0.078 | -2.12 | 0.27 | 0.136 | 1.75 | 0.49 |
| GB 02 TN | 0.85 | 0.061 | 0.23 | 0.021 | 3.15 | 0.73 |
| GB 04 TN | 0.84 | -0.78 | 0.21 | 0.329 | 1.54 | 0.44 |
| GB 00 TDN | -0.13 | -1.54 | 0.14 | 0.642 | 2.88 | 0.77 |

| | | | | | | |
|-----------|------|-------|------|-------|------|------|
| GB 01 TDN | 0.16 | -2.39 | 0.25 | 0.184 | 1.64 | 0.45 |
| GB 02 TDN | 0.72 | -0.66 | 0.24 | 0.016 | 3.06 | 0.71 |
| GB 04 TDN | 1.40 | 1.43 | 0.26 | 0.102 | 1.07 | 0.24 |

One Way Analysis of Variance

Data source: GB Water Quality Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: GB in Water Quality Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|-----------|----|---------|--------|--------|--------|
| GB 97 NOx | 16 | 0 | 0.0050 | 0.0050 | 0.0050 |
| GB 98 NOx | 15 | 0 | 0.0050 | 0.0050 | 0.0050 |
| GB 99 NOx | 13 | 0 | 0.0050 | 0.0050 | 0.0050 |
| GB 00 NOx | 18 | 0 | 0.0090 | 0.0050 | 0.045 |
| GB 01 NOx | 7 | 0 | 0.045 | 0.017 | 0.057 |
| GB 02 NOx | 16 | 0 | 0.0060 | 0.0050 | 0.018 |
| GB 04 NOx | 8 | 0 | 0.028 | 0.011 | 0.055 |

H = 42.54 with 6 degrees of freedom. (P = <0.001)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = <0.001)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|------------------------|---------------|------|--------|
| GB 01 NOx vs GB 98 NOx | 49.71 | 4.02 | Yes |
| GB 01 NOx vs GB 97 NOx | 47.40 | 3.88 | Yes |
| GB 01 NOx vs GB 99 NOx | 43.14 | 3.41 | Yes |
| GB 01 NOx vs GB 02 NOx | 26.93 | 2.20 | No |
| GB 01 NOx vs GB 00 NOx | 20.80 | 1.73 | No |
| GB 01 NOx vs GB 04 NOx | 9.90 | 0.71 | No |
| GB 04 NOx vs GB 98 NOx | 39.81 | 3.37 | Yes |
| GB 04 NOx vs GB 97 NOx | 37.50 | 3.21 | Yes |
| GB 04 NOx vs GB 99 NOx | 33.24 | 2.74 | No |
| GB 04 NOx vs GB 02 NOx | 17.03 | 1.46 | No |
| GB 04 NOx vs GB 00 NOx | 10.90 | 0.95 | No |
| GB 00 NOx vs GB 98 NOx | 28.92 | 3.06 | Yes |
| GB 00 NOx vs GB 97 NOx | 26.60 | 2.87 | No |
| GB 00 NOx vs GB 99 NOx | 22.34 | 2.27 | No |
| GB 00 NOx vs GB 02 NOx | 6.14 | 0.66 | No |
| GB 02 NOx vs GB 98 NOx | 22.78 | 2.35 | No |
| GB 02 NOx vs GB 97 NOx | 20.47 | 2.14 | No |
| GB 02 NOx vs GB 99 NOx | 16.20 | 1.61 | No |

| | | | |
|--|------|------|----|
| GB 99 NO _x vs GB 98 NO _x | 6.58 | 0.64 | No |
| GB 99 NO _x vs GB 97 NO _x | 4.26 | 0.42 | No |
| GB 97 NO _x vs GB 98 NO _x | 2.31 | 0.24 | No |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

One Way Analysis of Variance

Data source: GB Water Quality Trend Analysis

Normality Test: Failed (P = 0.003)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: GB in Water Quality Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|----------|----|---------|--------|-------|------|
| GB 00 TN | 12 | 0 | 0.27 | 0.19 | 0.34 |
| GB 01 TN | 7 | 0 | 0.20 | 0.19 | 0.35 |
| GB 02 TN | 16 | 0 | 0.16 | 0.14 | 0.27 |
| GB 04 TN | 8 | 0 | 0.14 | 0.075 | 0.31 |

H = 4.93 with 3 degrees of freedom. (P = 0.177)

The differences in the median values among the treatment groups are not great enough to exclude the possibility that the difference is due to random sampling variability; there is not a statistically significant difference (P = 0.177)

One Way Analysis of Variance

Data source: GB Water Quality Trend Analysis

Normality Test: Failed (P = 0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: GB in Water Quality Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|-----------|----|---------|--------|-------|------|
| GB 00 TDN | 12 | 0 | 0.25 | 0.16 | 0.32 |
| GB 01 TDN | 7 | 0 | 0.19 | 0.16 | 0.34 |
| GB 02 TDN | 16 | 0 | 0.15 | 0.13 | 0.28 |
| GB 04 TDN | 8 | 0 | 0.075 | 0.050 | 0.20 |

H = 6.92 with 3 degrees of freedom. (P = 0.074)

The differences in the median values among the treatment groups are not great enough to exclude the possibility that the difference is due to random sampling variability; there is not a statistically significant difference (P = 0.074)

Descriptive Statistics: Thursday, August 25, 2005, 09:55:47

Data source: GB in 5-Year Eelgrass Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|---------|------|---------|--------|---------|------------|--------------|
| GB 1999 | 12 | 0 | 498.67 | 127.42 | 36.78 | 80.96 |
| GB 2000 | 60 | 0 | 470.17 | 178.49 | 23.04 | 46.11 |
| GB 2001 | 60 | 0 | 372.83 | 123.95 | 16.00 | 32.02 |
| GB 2002 | 60 | 0 | 305.83 | 190.78 | 24.63 | 49.28 |
| GB 2004 | 60 | 0 | 300.17 | 204.78 | 26.44 | 52.90 |

| Column | Range | Max | Min | Median | 25% | 75% |
|---------|--------|--------|--------|--------|--------|--------|
| GB 1999 | 464.00 | 720.00 | 256.00 | 504.00 | 424.00 | 584.00 |
| GB 2000 | 820.00 | 950.00 | 130.00 | 465.00 | 340.00 | 580.00 |
| GB 2001 | 700.00 | 760.00 | 60.00 | 365.00 | 280.00 | 455.00 |
| GB 2002 | 670.00 | 670.00 | 0.00 | 340.00 | 160.00 | 410.00 |
| GB 2004 | 680.00 | 680.00 | 0.00 | 275.00 | 135.00 | 450.00 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|---------|----------|----------|-----------|-----------|----------|----------------|
| GB 1999 | -0.23 | 0.038 | 0.10 | 0.834 | 5984.00 | 3162624.00 |
| GB 2000 | 0.20 | -0.17 | 0.050 | 0.880 | 28210.00 | 15143100.00 |
| GB 2001 | 0.40 | 0.89 | 0.065 | 0.698 | 22370.00 | 9246700.00 |
| GB 2002 | -0.27 | -0.71 | 0.15 | 0.001 | 18350.00 | 7759500.00 |
| GB 2004 | 0.16 | -1.01 | 0.080 | 0.428 | 18010.00 | 7880100.00 |

One Way Analysis of Variance

Data source: GB Eelgrass Trend Analysis

Normality Test: Passed (P > 0.200)

Equal Variance Test: Failed (P = 0.002)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: GB in Eelgrass Trend analysis

| Group | N | Missing | Median | 25% | 75% |
|---------|----|---------|--------|--------|--------|
| GB 1999 | 12 | 0 | 504.00 | 424.00 | 584.00 |
| GB 2000 | 60 | 0 | 465.00 | 340.00 | 580.00 |
| GB 2001 | 60 | 0 | 365.00 | 280.00 | 455.00 |
| GB 2002 | 60 | 0 | 340.00 | 160.00 | 410.00 |
| GB 2004 | 60 | 0 | 275.00 | 135.00 | 450.00 |

H = 33.24 with 4 degrees of freedom. (P = <0.001)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = <0.001)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|--------------------|---------------|------|--------|
| GB 1999 vs GB 2004 | 77.77 | 3.37 | Yes |
| GB 1999 vs GB 2002 | 75.01 | 3.25 | Yes |
| GB 1999 vs GB 2001 | 54.87 | 2.38 | No |
| GB 1999 vs GB 2000 | 17.23 | 0.75 | No |
| GB 2000 vs GB 2004 | 60.53 | 4.55 | Yes |
| GB 2000 vs GB 2002 | 57.78 | 4.34 | Yes |
| GB 2000 vs GB 2001 | 37.63 | 2.83 | No |
| GB 2001 vs GB 2004 | 22.90 | 1.72 | No |
| GB 2001 vs GB 2002 | 20.14 | 1.51 | No |
| GB 2002 vs GB 2004 | 2.76 | 0.21 | No |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

Descriptive Statistics:

Data source: GB Algae Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean | |
|---------|------|---------|-------|---------|------------|--------------|--|
| GB 2000 | 24 | 0 | 25.83 | 21.40 | 4.37 | 9.04 | |
| GB 2001 | 60 | 0 | 14.18 | 15.53 | 2.01 | 4.01 | |
| GB 2002 | 60 | 0 | 23.37 | 25.29 | 3.26 | 6.53 | |
| GB 2004 | 60 | 0 | 21.30 | 22.52 | 2.91 | 5.82 | |

| Column | Range | Max | Min | Median | 25% | 75% |
|---------|--------|--------|------|--------|------|-------|
| GB 2000 | 75.00 | 75.00 | 0.00 | 22.50 | 7.50 | 50.00 |
| GB 2001 | 100.00 | 100.00 | 0.00 | 10.00 | 5.00 | 20.00 |
| GB 2002 | 90.00 | 90.00 | 0.00 | 10.00 | 5.00 | 45.00 |
| GB 2004 | 90.00 | 90.00 | 0.00 | 10.00 | 5.00 | 30.00 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|---------|----------|----------|-----------|-----------|---------|----------------|
| GB 2000 | 0.70 | -0.54 | 0.18 | 0.038 | 620.00 | 26550.00 |
| GB 2001 | 3.27 | 15.35 | 0.29 | <0.001 | 851.00 | 26301.00 |
| GB 2002 | 1.15 | 0.087 | 0.27 | <0.001 | 1402.00 | 70482.00 |
| GB 2004 | 1.41 | 1.35 | 0.26 | <0.001 | 1278.00 | 57140.00 |

One Way Analysis of Variance

Data source: GB Algae Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: GB in Algae Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|---------|----|---------|--------|------|-------|
| GB 2000 | 24 | 0 | 22.50 | 7.50 | 50.00 |
| GB 2001 | 60 | 0 | 10.00 | 5.00 | 20.00 |
| GB 2002 | 60 | 0 | 10.00 | 5.00 | 45.00 |
| GB 2004 | 60 | 0 | 10.00 | 5.00 | 30.00 |

H = 4.77 with 3 degrees of freedom. (P = 0.190)

The differences in the median values among the treatment groups are not great enough to exclude the possibility that the difference is due to random sampling variability; there is not a statistically significant difference ($P = 0.190$)

Appendix 8. The statistical reports generated for all of the analyses conducted in the synthesis of 2004 Long-Term Monitoring Report for Northwest Harbor.

Descriptive Statistics:

Data source: NWH Water Quality Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|------------|------|---------|--------|---------|------------|--------------|
| NWH 97 NOx | 47 | 0 | 0.0053 | 0.0020 | 0.00030 | 0.00060 |
| NWH 98 NOx | 42 | 0 | 0.0050 | 0.00015 | 0.000024 | 0.000048 |
| NWH 99 NOx | 36 | 0 | 0.0057 | 0.0021 | 0.00036 | 0.00073 |
| NWH 00 NOx | 21 | 0 | 0.016 | 0.024 | 0.0051 | 0.011 |
| NWH 01 NOx | 20 | 0 | 0.033 | 0.024 | 0.0054 | 0.011 |
| NWH 02 NOx | 18 | 0 | 0.027 | 0.036 | 0.0085 | 0.018 |
| NWH 04 NOx | 9 | 0 | 0.028 | 0.021 | 0.0069 | 0.016 |
| NWH 00 TN | 8 | 0 | 0.24 | 0.055 | 0.020 | 0.046 |
| NWH 01 TN | 20 | 0 | 0.21 | 0.084 | 0.019 | 0.039 |
| NWH 02 TN | 18 | 0 | 0.20 | 0.089 | 0.021 | 0.044 |
| NWH 04 TN | 9 | 0 | 0.18 | 0.19 | 0.063 | 0.15 |
| NWH 00 TDN | 8 | 0 | 0.24 | 0.050 | 0.018 | 0.042 |
| NWH 01 TDN | 20 | 0 | 0.20 | 0.080 | 0.018 | 0.038 |
| NWH 02 TDN | 18 | 0 | 0.19 | 0.083 | 0.019 | 0.041 |
| NWH 04 TDN | 9 | 0 | 0.17 | 0.19 | 0.063 | 0.15 |

| Column | Range | Max | Min | Median | 25% | 75% |
|------------|---------|--------|--------|--------|--------|--------|
| NWH 97 NOx | 0.014 | 0.019 | 0.0050 | 0.0050 | 0.0050 | 0.0050 |
| NWH 98 NOx | 0.00100 | 0.0060 | 0.0050 | 0.0050 | 0.0050 | 0.0050 |
| NWH 99 NOx | 0.0100 | 0.015 | 0.0050 | 0.0050 | 0.0050 | 0.0050 |
| NWH 00 NOx | 0.089 | 0.094 | 0.0050 | 0.0050 | 0.0050 | 0.011 |
| NWH 01 NOx | 0.075 | 0.080 | 0.0050 | 0.039 | 0.0075 | 0.051 |
| NWH 02 NOx | 0.10 | 0.11 | 0.0050 | 0.0075 | 0.0050 | 0.034 |
| NWH 04 NOx | 0.056 | 0.061 | 0.0050 | 0.018 | 0.014 | 0.047 |
| NWH 00 TN | 0.18 | 0.32 | 0.14 | 0.23 | 0.22 | 0.28 |
| NWH 01 TN | 0.25 | 0.36 | 0.11 | 0.17 | 0.16 | 0.26 |
| NWH 02 TN | 0.32 | 0.40 | 0.080 | 0.19 | 0.13 | 0.28 |
| NWH 04 TN | 0.56 | 0.61 | 0.050 | 0.080 | 0.050 | 0.24 |
| NWH 00 TDN | 0.15 | 0.31 | 0.16 | 0.23 | 0.21 | 0.29 |
| NWH 01 TDN | 0.25 | 0.36 | 0.11 | 0.17 | 0.14 | 0.24 |
| NWH 02 TDN | 0.29 | 0.37 | 0.080 | 0.16 | 0.12 | 0.27 |
| NWH 04 TDN | 0.53 | 0.58 | 0.050 | 0.060 | 0.050 | 0.25 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|------------|----------|----------|-----------|-----------|------|----------------|
| NWH 97 NOx | 6.80 | 46.49 | 0.52 | <0.001 | 0.25 | 0.0015 |
| NWH 98 NOx | 6.48 | 42.00 | 0.54 | <0.001 | 0.21 | 0.0011 |
| NWH 99 NOx | 3.44 | 11.73 | 0.49 | <0.001 | 0.21 | 0.0013 |
| NWH 00 NOx | 2.58 | 6.18 | 0.39 | <0.001 | 0.33 | 0.016 |
| NWH 01 NOx | 0.20 | -1.23 | 0.19 | 0.057 | 0.66 | 0.033 |
| NWH 02 NOx | 1.45 | 0.50 | 0.35 | <0.001 | 0.49 | 0.035 |
| NWH 04 NOx | 0.84 | -0.98 | 0.26 | 0.080 | 0.25 | 0.010 |
| NWH 00 TN | -0.16 | 0.62 | 0.19 | 0.513 | 1.90 | 0.47 |
| NWH 01 TN | 0.93 | -0.53 | 0.25 | 0.002 | 4.19 | 1.01 |
| NWH 02 TN | 0.65 | -0.35 | 0.14 | 0.386 | 3.63 | 0.87 |
| NWH 04 TN | 1.82 | 3.21 | 0.25 | 0.099 | 1.58 | 0.56 |

| | | | | | | |
|------------|-------|-------|------|--------|------|------|
| NWH 00 TDN | -0.14 | -0.57 | 0.17 | 0.603 | 1.94 | 0.49 |
| NWH 01 TDN | 1.12 | -0.20 | 0.31 | <0.001 | 4.03 | 0.94 |
| NWH 02 TDN | 0.66 | -0.46 | 0.17 | 0.187 | 3.35 | 0.74 |
| NWH 04 TDN | 1.66 | 1.96 | 0.31 | 0.014 | 1.52 | 0.54 |

One Way Analysis of Variance

Data source: NWH Water Quality Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: NWH in Water Quality Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|------------|----|---------|--------|--------|--------|
| NWH 97 NOx | 47 | 0 | 0.0050 | 0.0050 | 0.0050 |
| NWH 98 NOx | 42 | 0 | 0.0050 | 0.0050 | 0.0050 |
| NWH 99 NOx | 36 | 0 | 0.0050 | 0.0050 | 0.0050 |
| NWH 00 NOx | 21 | 0 | 0.0050 | 0.0050 | 0.011 |
| NWH 01 NOx | 20 | 0 | 0.039 | 0.0075 | 0.051 |
| NWH 02 NOx | 18 | 0 | 0.0075 | 0.0050 | 0.034 |
| NWH 04 NOx | 9 | 0 | 0.018 | 0.014 | 0.047 |

H = 92.31 with 6 degrees of freedom. (P = <0.001)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = <0.001)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|--------------------------|---------------|------|--------|
| NWH 04 NOx vs NWH 98 NOx | 87.55 | 4.27 | Yes |
| NWH 04 NOx vs NWH 97 NOx | 85.65 | 4.21 | Yes |
| NWH 04 NOx vs NWH 99 NOx | 77.69 | 3.73 | Yes |
| NWH 04 NOx vs NWH 00 NOx | 45.04 | 2.02 | No |
| NWH 04 NOx vs NWH 02 NOx | 33.83 | 1.48 | No |
| NWH 04 NOx vs NWH 01 NOx | 7.90 | 0.35 | No |
| NWH 01 NOx vs NWH 98 NOx | 79.65 | 5.25 | Yes |
| NWH 01 NOx vs NWH 97 NOx | 77.75 | 5.21 | Yes |
| NWH 01 NOx vs NWH 99 NOx | 69.79 | 4.48 | Yes |
| NWH 01 NOx vs NWH 00 NOx | 37.14 | 2.13 | No |
| NWH 01 NOx vs NWH 02 NOx | 25.93 | 1.43 | No |
| NWH 02 NOx vs NWH 98 NOx | 53.72 | 3.41 | Yes |
| NWH 02 NOx vs NWH 97 NOx | 51.82 | 3.35 | Yes |
| NWH 02 NOx vs NWH 99 NOx | 43.86 | 2.72 | No |
| NWH 02 NOx vs NWH 00 NOx | 11.21 | 0.62 | No |
| NWH 00 NOx vs NWH 98 NOx | 42.51 | 2.85 | No |

| | | | |
|--------------------------|-------|------|----|
| NWH 00 NOx vs NWH 97 NOx | 40.61 | 2.77 | No |
| NWH 00 NOx vs NWH 99 NOx | 32.65 | 2.13 | No |
| NWH 99 NOx vs NWH 98 NOx | 9.86 | 0.78 | No |
| NWH 99 NOx vs NWH 97 NOx | 7.96 | 0.64 | No |
| NWH 97 NOx vs NWH 98 NOx | 1.90 | 0.16 | No |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

One Way Analysis of Variance

Data source: NWH Water Quality Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: NWH in Water Quality Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|-----------|----|---------|--------|-------|------|
| NWH 00 TN | 8 | 0 | 0.23 | 0.22 | 0.28 |
| NWH 01 TN | 20 | 0 | 0.17 | 0.16 | 0.26 |
| NWH 02 TN | 18 | 0 | 0.19 | 0.13 | 0.28 |
| NWH 04 TN | 9 | 0 | 0.080 | 0.050 | 0.24 |

H = 5.46 with 3 degrees of freedom. (P = 0.141)

The differences in the median values among the treatment groups are not great enough to exclude the possibility that the difference is due to random sampling variability; there is not a statistically significant difference (P = 0.141)

One Way Analysis of Variance

Data source: NWH Water Quality Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: NWH in Water Quality Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|------------|----|---------|--------|-------|------|
| NWH 00 TDN | 8 | 0 | 0.23 | 0.21 | 0.29 |
| NWH 01 TDN | 20 | 0 | 0.17 | 0.14 | 0.24 |
| NWH 02 TDN | 18 | 0 | 0.16 | 0.12 | 0.27 |
| NWH 04 TDN | 9 | 0 | 0.060 | 0.050 | 0.25 |

H = 7.40 with 3 degrees of freedom. (P = 0.060)

The differences in the median values among the treatment groups are not great enough to exclude the possibility that the difference is due to random sampling variability; there is not a statistically significant difference (P = 0.060)

Descriptive Statistics:

Data source: NWH Eelgrass Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|----------|------|---------|--------|---------|------------|--------------|
| NWH 1997 | 3 | 0 | 209.33 | 41.05 | 23.70 | 101.98 |
| NWH 1998 | 12 | 0 | 310.67 | 72.67 | 20.98 | 46.17 |
| NWH 1999 | 12 | 0 | 506.67 | 196.71 | 56.79 | 124.98 |
| NWH 2000 | 60 | 0 | 329.83 | 166.03 | 21.43 | 42.89 |
| NWH 2001 | 60 | 0 | 408.83 | 155.71 | 20.10 | 40.22 |
| NWH 2002 | 60 | 0 | 349.83 | 146.15 | 18.87 | 37.76 |
| NWH 2004 | 60 | 0 | 290.50 | 141.81 | 18.31 | 36.63 |

| Column | Range | Max | Min | Median | 25% | 75% |
|----------|--------|--------|--------|--------|--------|--------|
| NWH 1997 | 80.00 | 244.00 | 164.00 | 220.00 | 178.00 | 238.00 |
| NWH 1998 | 240.00 | 400.00 | 160.00 | 336.00 | 272.00 | 360.00 |
| NWH 1999 | 704.00 | 864.00 | 160.00 | 520.00 | 368.00 | 616.00 |
| NWH 2000 | 640.00 | 720.00 | 80.00 | 320.00 | 185.00 | 475.00 |
| NWH 2001 | 700.00 | 820.00 | 120.00 | 400.00 | 280.00 | 520.00 |
| NWH 2002 | 730.00 | 800.00 | 70.00 | 330.00 | 245.00 | 445.00 |
| NWH 2004 | 650.00 | 650.00 | 0.00 | 300.00 | 190.00 | 390.00 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|----------|----------|----------|-----------|-----------|----------|----------------|
| NWH 1997 | -1.09 | -- | 0.27 | 0.429 | 628.00 | 134832.00 |
| NWH 1998 | -1.01 | 0.18 | 0.22 | 0.128 | 3728.00 | 1216256.00 |
| NWH 1999 | -0.060 | -0.024 | 0.15 | 0.613 | 6080.00 | 3506176.00 |
| NWH 2000 | 0.38 | -0.77 | 0.13 | 0.010 | 19790.00 | 8153700.00 |
| NWH 2001 | 0.31 | 0.041 | 0.067 | 0.664 | 24530.00 | 11459100.00 |
| NWH 2002 | 0.71 | 0.86 | 0.087 | 0.297 | 20990.00 | 8603300.00 |
| NWH 2004 | 0.18 | -0.097 | 0.072 | 0.562 | 17430.00 | 6249900.00 |

One Way Analysis of Variance

Data source: NWH Eelgrass Trend Analysis

Normality Test: Passed (P = 0.197)

Equal Variance Test: Passed (P = 0.068)

| Group Name | N | Missing | Mean | Std Dev | SEM |
|------------|----|---------|--------|---------|-------|
| NWH 1997 | 3 | 0 | 209.33 | 41.05 | 23.70 |
| NWH 1998 | 12 | 0 | 310.67 | 72.67 | 20.98 |
| NWH 1999 | 12 | 0 | 506.67 | 196.71 | 56.79 |
| NWH 2000 | 60 | 0 | 329.83 | 166.03 | 21.43 |
| NWH 2001 | 60 | 0 | 408.83 | 155.71 | 20.10 |
| NWH 2002 | 60 | 0 | 349.83 | 146.15 | 18.87 |
| NWH 2004 | 60 | 0 | 290.50 | 141.81 | 18.31 |

| Source of Variation | DF | SS | MS | F | P |
|---------------------|-----|------------|-----------|------|--------|
| Between Groups | 6 | 816678.82 | 136113.14 | 5.91 | <0.001 |
| Residual | 260 | 5990604.00 | 23040.78 | | |

Total 266 6807282.82

The differences in the mean values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = <0.001).

Power of performed test with alpha = 0.05: 0.994

All Pairwise Multiple Comparison Procedures (Tukey Test):

Comparisons for factor:

| Comparison | Diff of Means | p | q | P | P<0.050 |
|-----------------------|---------------|---|------|--------|---------|
| NWH 1999 vs. NWH 1997 | 297.33 | 7 | 4.29 | 0.039 | Yes |
| NWH 1999 vs. NWH 2004 | 216.17 | 7 | 6.37 | <0.001 | Yes |
| NWH 1999 vs. NWH 1998 | 196.00 | 7 | 4.47 | 0.026 | Yes |
| NWH 1999 vs. NWH 2000 | 176.83 | 7 | 5.21 | 0.004 | Yes |
| NWH 1999 vs. NWH 2002 | 156.83 | 7 | 4.62 | 0.019 | Yes |
| NWH 1999 vs. NWH 2001 | 97.83 | 7 | 2.88 | 0.390 | No |
| NWH 2001 vs. NWH 1997 | 199.50 | 7 | 3.14 | 0.284 | No |
| NWH 2001 vs. NWH 2004 | 118.33 | 7 | 6.04 | <0.001 | No |
| NWH 2001 vs. NWH 1998 | 98.17 | 7 | 2.89 | 0.386 | No |
| NWH 2001 vs. NWH 2000 | 79.00 | 7 | 4.03 | 0.066 | No |
| NWH 2001 vs. NWH 2002 | 59.00 | 7 | 3.01 | 0.335 | No |
| NWH 2002 vs. NWH 1997 | 140.50 | 7 | 2.21 | 0.705 | No |
| NWH 2002 vs. NWH 2004 | 59.33 | 7 | 3.03 | 0.328 | No |
| NWH 2002 vs. NWH 1998 | 39.17 | 7 | 1.15 | 0.983 | No |
| NWH 2002 vs. NWH 2000 | 20.00 | 7 | 1.02 | 0.991 | No |
| NWH 2000 vs. NWH 1997 | 120.50 | 7 | 1.90 | 0.832 | No |
| NWH 2000 vs. NWH 2004 | 39.33 | 7 | 2.01 | 0.792 | No |
| NWH 2000 vs. NWH 1998 | 19.17 | 7 | 0.56 | 1.000 | No |
| NWH 1998 vs. NWH 1997 | 101.33 | 7 | 1.46 | 0.946 | No |
| NWH 1998 vs. NWH 2004 | 20.17 | 7 | 0.59 | 1.000 | No |
| NWH 2004 vs. NWH 1997 | 81.17 | 7 | 1.28 | 0.972 | No |

Descriptive Statistics:

Data source: NWH Algae Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|----------|------|---------|-------|---------|------------|--------------|
| NWH 2000 | 24 | 0 | 38.29 | 28.73 | 5.87 | 12.13 |
| NWH 2001 | 60 | 0 | 30.92 | 23.91 | 3.09 | 6.18 |
| NWH 2002 | 60 | 0 | 64.25 | 29.38 | 3.79 | 7.59 |
| NWH 2004 | 60 | 0 | 81.42 | 24.06 | 3.11 | 6.22 |

| Column | Range | Max | Min | Median | 25% | 75% |
|----------|--------|--------|-------|--------|-------|--------|
| NWH 2000 | 89.00 | 90.00 | 1.00 | 40.00 | 10.00 | 60.00 |
| NWH 2001 | 100.00 | 100.00 | 0.00 | 30.00 | 10.00 | 37.50 |
| NWH 2002 | 90.00 | 100.00 | 10.00 | 75.00 | 40.00 | 90.00 |
| NWH 2004 | 80.00 | 100.00 | 20.00 | 92.50 | 75.00 | 100.00 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|----------|----------|----------|-----------|-----------|---------|----------------|
| NWH 2000 | 0.23 | -1.03 | 0.14 | 0.284 | 919.00 | 54179.00 |
| NWH 2001 | 1.33 | 1.51 | 0.25 | <0.001 | 1855.00 | 91075.00 |

| | | | | | | |
|----------|-------|-------|------|--------|---------|-----------|
| NWH 2002 | -0.44 | -1.11 | 0.24 | <0.001 | 3855.00 | 298625.00 |
| NWH 2004 | -1.25 | 0.38 | 0.26 | <0.001 | 4885.00 | 431875.00 |

One Way Analysis of Variance

Data source: NWH Algae Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: NWH in Algae Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|----------|----|---------|--------|-------|--------|
| NWH 2000 | 24 | 0 | 40.00 | 10.00 | 60.00 |
| NWH 2001 | 60 | 0 | 30.00 | 10.00 | 37.50 |
| NWH 2002 | 60 | 0 | 75.00 | 40.00 | 90.00 |
| NWH 2004 | 60 | 0 | 92.50 | 75.00 | 100.00 |

H = 75.95 with 3 degrees of freedom. (P = <0.001)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = <0.001)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|----------------------|---------------|------|--------|
| NWH 2004 vs NWH 2001 | 86.82 | 8.06 | Yes |
| NWH 2004 vs NWH 2000 | 77.17 | 5.41 | Yes |
| NWH 2004 vs NWH 2002 | 32.05 | 2.97 | Yes |
| NWH 2002 vs NWH 2001 | 54.77 | 5.08 | Yes |
| NWH 2002 vs NWH 2000 | 45.12 | 3.16 | Yes |
| NWH 2000 vs NWH 2001 | 9.66 | 0.68 | No |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

Appendix 9. The statistical reports generated for all of the analyses conducted in the synthesis of 2004 Long-Term Monitoring Report for Orient Harbor.

Descriptive Statistics:

Data source: OH Water Quality Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|-----------|------|---------|--------|---------|------------|--------------|
| OH 97 NOx | 46 | 0 | 0.0054 | 0.0020 | 0.00030 | 0.00060 |
| OH 98 NOx | 70 | 0 | 0.0061 | 0.0045 | 0.00054 | 0.0011 |
| OH 99 NOx | 25 | 0 | 0.0069 | 0.0039 | 0.00077 | 0.0016 |
| OH 00 NOx | 21 | 0 | 0.012 | 0.012 | 0.0027 | 0.0055 |
| OH 01 NOx | 20 | 0 | 0.038 | 0.030 | 0.0068 | 0.014 |
| OH 02 NOx | 17 | 0 | 0.021 | 0.022 | 0.0053 | 0.011 |
| OH 04 NOx | 9 | 0 | 0.024 | 0.031 | 0.010 | 0.024 |
| OH 00 TN | 8 | 0 | 0.25 | 0.062 | 0.022 | 0.052 |
| OH 01 TN | 20 | 0 | 0.21 | 0.085 | 0.019 | 0.040 |
| OH 02 TN | 17 | 0 | 0.19 | 0.090 | 0.022 | 0.046 |
| OH 04 TN | 9 | 0 | 0.14 | 0.12 | 0.039 | 0.090 |
| OH 00 TDN | 8 | 0 | 0.25 | 0.059 | 0.021 | 0.050 |
| OH 01 TDN | 20 | 0 | 0.20 | 0.079 | 0.018 | 0.037 |
| OH 02 TDN | 17 | 0 | 0.19 | 0.082 | 0.020 | 0.042 |
| OH 04 TDN | 9 | 1 | 0.14 | 0.14 | 0.048 | 0.11 |

| Column | Range | Max | Min | Median | 25% | 75% |
|-----------|-------|-------|--------|--------|--------|--------|
| OH 97 NOx | 0.013 | 0.018 | 0.0050 | 0.0050 | 0.0050 | 0.0050 |
| OH 98 NOx | 0.029 | 0.034 | 0.0050 | 0.0050 | 0.0050 | 0.0050 |
| OH 99 NOx | 0.013 | 0.018 | 0.0050 | 0.0050 | 0.0050 | 0.0073 |
| OH 00 NOx | 0.050 | 0.055 | 0.0050 | 0.0050 | 0.0050 | 0.014 |
| OH 01 NOx | 0.097 | 0.10 | 0.0050 | 0.029 | 0.013 | 0.057 |
| OH 02 NOx | 0.081 | 0.086 | 0.0050 | 0.011 | 0.0050 | 0.028 |
| OH 04 NOx | 0.074 | 0.079 | 0.0050 | 0.0100 | 0.0050 | 0.030 |
| OH 00 TN | 0.18 | 0.33 | 0.15 | 0.26 | 0.21 | 0.31 |
| OH 01 TN | 0.26 | 0.37 | 0.11 | 0.19 | 0.14 | 0.27 |
| OH 02 TN | 0.31 | 0.39 | 0.080 | 0.19 | 0.12 | 0.25 |
| OH 04 TN | 0.32 | 0.37 | 0.050 | 0.070 | 0.050 | 0.20 |
| OH 00 TDN | 0.16 | 0.33 | 0.17 | 0.24 | 0.21 | 0.30 |
| OH 01 TDN | 0.24 | 0.35 | 0.11 | 0.17 | 0.14 | 0.26 |
| OH 02 TDN | 0.29 | 0.38 | 0.090 | 0.17 | 0.11 | 0.24 |
| OH 04 TDN | 0.39 | 0.44 | 0.050 | 0.070 | 0.050 | 0.19 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|-----------|----------|----------|-----------|-----------|------|----------------|
| OH 97 NOx | 5.87 | 36.39 | 0.52 | <0.001 | 0.25 | 0.0015 |
| OH 98 NOx | 4.98 | 26.24 | 0.48 | <0.001 | 0.43 | 0.0040 |
| OH 99 NOx | 2.26 | 4.32 | 0.41 | <0.001 | 0.17 | 0.0015 |
| OH 00 NOx | 2.58 | 7.52 | 0.28 | <0.001 | 0.26 | 0.0061 |
| OH 01 NOx | 0.81 | -0.51 | 0.17 | 0.111 | 0.76 | 0.046 |
| OH 02 NOx | 1.97 | 4.03 | 0.27 | 0.002 | 0.35 | 0.015 |
| OH 04 NOx | 1.56 | 0.63 | 0.40 | <0.001 | 0.21 | 0.013 |
| OH 00 TN | -0.27 | -0.55 | 0.14 | 0.757 | 2.03 | 0.54 |
| OH 01 TN | 0.85 | -0.62 | 0.16 | 0.178 | 4.24 | 1.03 |
| OH 02 TN | 0.67 | -0.25 | 0.18 | 0.176 | 3.31 | 0.77 |
| OH 04 TN | 1.08 | 0.12 | 0.28 | 0.036 | 1.27 | 0.29 |
| OH 00 TDN | 0.29 | -0.78 | 0.18 | 0.561 | 1.98 | 0.52 |

| | | | | | | |
|-----------|------|-------|------|-------|------|------|
| OH 01 TDN | 0.87 | -0.70 | 0.19 | 0.070 | 3.94 | 0.89 |
| OH 02 TDN | 0.87 | 0.25 | 0.16 | 0.266 | 3.15 | 0.69 |
| OH 04 TDN | 1.93 | 3.83 | 0.29 | 0.048 | 1.10 | 0.28 |

One Way Analysis of Variance

Data source: OH Water Quality Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: OH Water Quality Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|-----------|----|---------|---------|---------|---------|
| OH 97 NOx | 46 | 0 | 0.00500 | 0.00500 | 0.00500 |
| OH 98 NOx | 70 | 0 | 0.00500 | 0.00500 | 0.00500 |
| OH 99 NOx | 25 | 0 | 0.00500 | 0.00500 | 0.00725 |
| OH 00 NOx | 21 | 0 | 0.00500 | 0.00500 | 0.0143 |
| OH 01 NOx | 20 | 0 | 0.0290 | 0.0130 | 0.0575 |
| OH 02 NOx | 17 | 0 | 0.0110 | 0.00500 | 0.0283 |
| OH 04 NOx | 9 | 0 | 0.01000 | 0.00500 | 0.0298 |

H = 87.552 with 6 degrees of freedom. (P = <0.001)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = <0.001)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|------------------------|---------------|-------|--------|
| OH 01 NOx vs OH 97 NOx | 97.188 | 6.029 | Yes |
| OH 01 NOx vs OH 98 NOx | 92.411 | 6.056 | Yes |
| OH 01 NOx vs OH 99 NOx | 77.565 | 4.296 | Yes |
| OH 01 NOx vs OH 00 NOx | 54.025 | 2.873 | No |
| OH 01 NOx vs OH 04 NOx | 42.192 | 1.746 | No |
| OH 01 NOx vs OH 02 NOx | 28.496 | 1.435 | No |
| OH 02 NOx vs OH 97 NOx | 68.692 | 4.021 | Yes |
| OH 02 NOx vs OH 98 NOx | 63.915 | 3.927 | Yes |
| OH 02 NOx vs OH 99 NOx | 49.069 | 2.593 | No |
| OH 02 NOx vs OH 00 NOx | 25.529 | 1.300 | No |
| OH 02 NOx vs OH 04 NOx | 13.696 | 0.552 | No |
| OH 04 NOx vs OH 97 NOx | 54.996 | 2.507 | No |
| OH 04 NOx vs OH 98 NOx | 50.219 | 2.356 | No |
| OH 04 NOx vs OH 99 NOx | 35.373 | 1.512 | No |
| OH 04 NOx vs OH 00 NOx | 11.833 | 0.493 | No |
| OH 00 NOx vs OH 97 NOx | 43.163 | 2.723 | No |
| OH 00 NOx vs OH 98 NOx | 38.386 | 2.563 | No |

| | | | |
|--|--------|-------|----|
| OH 00 NO _x vs OH 99 NO _x | 23.540 | 1.321 | No |
| OH 99 NO _x vs OH 97 NO _x | 19.623 | 1.312 | No |
| OH 99 NO _x vs OH 98 NO _x | 14.846 | 1.059 | No |
| OH 98 NO _x vs OH 97 NO _x | 4.777 | 0.418 | No |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

One Way Analysis of Variance

Data source: OH Water Quality Trend Analysis

Normality Test: Passed (P = 0.032)

Equal Variance Test: Passed (P = 0.388)

| Group Name | N | Missing | Mean | Std Dev | SEM |
|------------|----|---------|------|---------|-------|
| OH 00 TN | 8 | 0 | 0.25 | 0.062 | 0.022 |
| OH 01 TN | 20 | 0 | 0.21 | 0.085 | 0.019 |
| OH 02 TN | 17 | 0 | 0.19 | 0.090 | 0.022 |
| OH 04 TN | 9 | 0 | 0.14 | 0.12 | 0.039 |

| Source of Variation | DF | SS | MS | F | P |
|---------------------|----|-------|--------|------|-------|
| Between Groups | 3 | 0.057 | 0.019 | 2.36 | 0.083 |
| Residual | 50 | 0.40 | 0.0081 | | |
| Total | 53 | 0.46 | | | |

The differences in the mean values among the treatment groups are not great enough to exclude the possibility that the difference is due to random sampling variability; there is not a statistically significant difference (P = 0.083).

Power of performed test with alpha = 0.05: 0.330

The power of the performed test (0.330) is below the desired power of 0.800.

You should interpret the negative findings cautiously.

One Way Analysis of Variance

Data source: OH Water Quality Trend Analysis

Normality Test: Failed (P = 0.003)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: OH in Water Quality Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|-----------|----|---------|--------|-------|------|
| OH 00 TDN | 8 | 0 | 0.24 | 0.21 | 0.30 |
| OH 01 TDN | 20 | 0 | 0.17 | 0.14 | 0.26 |
| OH 02 TDN | 17 | 0 | 0.17 | 0.11 | 0.24 |
| OH 04 TDN | 9 | 1 | 0.070 | 0.050 | 0.19 |

H = 8.92 with 3 degrees of freedom. (P = 0.030)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = 0.030)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|------------------------|---------------|------|--------|
| OH 00 TDN vs OH 04 TDN | 22.69 | 2.94 | Yes |
| OH 00 TDN vs OH 02 TDN | 12.97 | 1.96 | No |
| OH 00 TDN vs OH 01 TDN | 10.38 | 1.61 | No |
| OH 01 TDN vs OH 04 TDN | 12.31 | 1.91 | No |
| OH 01 TDN vs OH 02 TDN | 2.60 | 0.51 | No |
| OH 02 TDN vs OH 04 TDN | 9.72 | 1.47 | No |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

Descriptive Statistics:

Data source: OH in 5-Year Eelgrass Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|---------|------|---------|--------|---------|------------|--------------|
| OH 1997 | 3 | 0 | 573.33 | 118.28 | 68.29 | 293.82 |
| OH 1998 | 10 | 0 | 696.00 | 260.00 | 82.22 | 185.99 |
| OH 1999 | 12 | 0 | 586.67 | 171.29 | 49.45 | 108.83 |
| OH 2000 | 60 | 0 | 487.83 | 200.57 | 25.89 | 51.81 |
| OH 2001 | 60 | 0 | 451.50 | 127.24 | 16.43 | 32.87 |
| OH 2002 | 60 | 0 | 229.50 | 103.77 | 13.40 | 26.81 |
| OH 2004 | 60 | 0 | 55.50 | 113.26 | 14.62 | 29.26 |

| Column | Range | Max | Min | Median | 25% | 75% |
|---------|--------|---------|--------|--------|--------|--------|
| OH 1997 | 236.00 | 696.00 | 460.00 | 564.00 | 486.00 | 663.00 |
| OH 1998 | 880.00 | 1088.00 | 208.00 | 712.00 | 576.00 | 832.00 |
| OH 1999 | 496.00 | 832.00 | 336.00 | 600.00 | 456.00 | 720.00 |
| OH 2000 | 950.00 | 990.00 | 40.00 | 460.00 | 350.00 | 610.00 |
| OH 2001 | 720.00 | 780.00 | 60.00 | 460.00 | 360.00 | 540.00 |
| OH 2002 | 430.00 | 440.00 | 10.00 | 235.00 | 150.00 | 305.00 |
| OH 2004 | 650.00 | 650.00 | 0.00 | 0.00 | 0.00 | 75.00 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|---------|----------|----------|-----------|-----------|----------|----------------|
| OH 1997 | 0.35 | -- | 0.20 | 0.633 | 1720.00 | 1014112.00 |
| OH 1998 | -0.52 | 0.22 | 0.16 | 0.560 | 6960.00 | 5452544.00 |
| OH 1999 | -0.12 | -1.26 | 0.16 | 0.514 | 7040.00 | 4452864.00 |
| OH 2000 | 0.34 | -0.074 | 0.089 | 0.278 | 29270.00 | 16652300.00 |
| OH 2001 | -0.100 | 0.72 | 0.064 | 0.708 | 27090.00 | 13186300.00 |
| OH 2002 | -0.065 | -0.75 | 0.100 | 0.139 | 13770.00 | 3795500.00 |
| OH 2004 | 3.15 | 12.58 | 0.32 | <0.001 | 3330.00 | 941700.00 |

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: OH Eelgrass Trend Analysis

Normality Test: Failed (P = <0.001)

| Group | N | Missing | Median | 25% | 75% |
|---------|----|---------|--------|--------|--------|
| OH 1997 | 3 | 0 | 564.00 | 486.00 | 663.00 |
| OH 1998 | 10 | 0 | 712.00 | 576.00 | 832.00 |
| OH 1999 | 12 | 0 | 600.00 | 456.00 | 720.00 |
| OH 2000 | 60 | 0 | 460.00 | 350.00 | 610.00 |
| OH 2001 | 60 | 0 | 460.00 | 360.00 | 540.00 |
| OH 2002 | 60 | 0 | 235.00 | 150.00 | 305.00 |
| OH 2004 | 60 | 0 | 0.00 | 0.00 | 75.00 |

H = 177.13 with 6 degrees of freedom. (P = <0.001)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = <0.001)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|--------------------|---------------|-------|--------|
| OH 1998 vs OH 2004 | 184.23 | 7.04 | Yes |
| OH 1998 vs OH 2002 | 126.71 | 4.84 | Yes |
| OH 1998 vs OH 2001 | 46.41 | 1.77 | No |
| OH 1998 vs OH 2000 | 42.40 | 1.62 | No |
| OH 1998 vs OH 1999 | 12.36 | 0.38 | No |
| OH 1998 vs OH 1997 | 7.15 | 0.14 | No |
| OH 1997 vs OH 2004 | 177.08 | 3.91 | Yes |
| OH 1997 vs OH 2002 | 119.56 | 2.64 | No |
| OH 1997 vs OH 2001 | 39.26 | 0.87 | No |
| OH 1997 vs OH 2000 | 35.25 | 0.78 | No |
| OH 1997 vs OH 1999 | 5.21 | 0.11 | No |
| OH 1999 vs OH 2004 | 171.88 | 7.09 | Yes |
| OH 1999 vs OH 2002 | 114.35 | 4.72 | No |
| OH 1999 vs OH 2001 | 34.05 | 1.40 | No |
| OH 1999 vs OH 2000 | 30.04 | 1.24 | No |
| OH 2000 vs OH 2004 | 141.83 | 10.14 | Yes |
| OH 2000 vs OH 2002 | 84.31 | 6.03 | No |
| OH 2000 vs OH 2001 | 4.01 | 0.29 | No |
| OH 2001 vs OH 2004 | 137.83 | 9.85 | Yes |
| OH 2001 vs OH 2002 | 80.30 | 5.74 | No |
| OH 2002 vs OH 2004 | 57.52 | 4.11 | Yes |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

Descriptive Statistics:

Data source: OH Algae Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|---------|------|---------|-------|---------|------------|--------------|
| OH 2000 | 24 | 0 | 7.13 | 11.95 | 2.44 | 5.04 |
| OH 2001 | 60 | 0 | 37.08 | 23.80 | 3.07 | 6.15 |
| OH 2002 | 60 | 0 | 12.70 | 16.46 | 2.12 | 4.25 |
| OH 2004 | 60 | 0 | 1.13 | 3.66 | 0.47 | 0.95 |

| Column | Range | Max | Min | Median | 25% | 75% |
|---------|-------|--------|------|--------|-------|-------|
| OH 2000 | 50.00 | 50.00 | 0.00 | 3.00 | 0.00 | 7.50 |
| OH 2001 | 95.00 | 100.00 | 5.00 | 30.00 | 20.00 | 50.00 |
| OH 2002 | 90.00 | 90.00 | 0.00 | 7.50 | 5.00 | 10.00 |
| OH 2004 | 25.00 | 25.00 | 0.00 | 0.00 | 0.00 | 1.00 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|---------|----------|----------|-----------|-----------|---------|----------------|
| OH 2000 | 2.45 | 6.66 | 0.32 | <0.001 | 171.00 | 4501.00 |
| OH 2001 | 0.82 | 0.098 | 0.15 | 0.002 | 2225.00 | 115925.00 |
| OH 2002 | 2.61 | 8.12 | 0.35 | <0.001 | 762.00 | 25662.00 |
| OH 2004 | 5.36 | 32.04 | 0.45 | <0.001 | 68.00 | 868.00 |

One Way Analysis of Variance

Data source: OH Algae Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: OH in Algae Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|---------|----|---------|--------|-------|-------|
| OH 2000 | 24 | 0 | 3.00 | 0.00 | 7.50 |
| OH 2001 | 60 | 0 | 30.00 | 20.00 | 50.00 |
| OH 2002 | 60 | 0 | 7.50 | 5.00 | 10.00 |
| OH 2004 | 60 | 0 | 0.00 | 0.00 | 1.00 |

H = 130.69 with 3 degrees of freedom. (P = <0.001)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = <0.001)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|--------------------|---------------|-------|--------|
| OH 2001 vs OH 2004 | 118.43 | 10.99 | Yes |
| OH 2001 vs OH 2000 | 87.77 | 6.16 | Yes |
| OH 2001 vs OH 2002 | 50.80 | 4.71 | Yes |
| OH 2002 vs OH 2004 | 67.63 | 6.28 | Yes |
| OH 2002 vs OH 2000 | 36.97 | 2.59 | No |
| OH 2000 vs OH 2004 | 30.67 | 2.15 | No |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

Appendix 10. The statistical reports generated for all of the analyses conducted in the synthesis of 2004 Long-Term Monitoring Report for Southold Harbor.

Descriptive Statistics:

Data source: SB Water Quality Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|-----------|------|---------|--------|---------|------------|--------------|
| SB 97 NOx | 24 | 0 | 0.0059 | 0.0028 | 0.00056 | 0.0012 |
| SB 98 NOx | 15 | 0 | 0.0047 | 0.0010 | 0.00027 | 0.00057 |
| SB 99 NOx | 15 | 0 | 0.0094 | 0.0069 | 0.0018 | 0.0038 |
| SB 00 NOx | 11 | 0 | 0.019 | 0.021 | 0.0063 | 0.014 |
| SB 01 NOx | 13 | 0 | 0.024 | 0.023 | 0.0064 | 0.014 |
| SB 02 NOx | 11 | 0 | 0.023 | 0.016 | 0.0050 | 0.011 |
| SB 04 NOx | 4 | 0 | 0.025 | 0.017 | 0.0086 | 0.027 |
| SB 00 TN | 5 | 0 | 0.25 | 0.053 | 0.024 | 0.065 |
| SB 01 TN | 13 | 0 | 0.22 | 0.086 | 0.024 | 0.052 |
| SB 02 TN | 11 | 0 | 0.19 | 0.086 | 0.026 | 0.058 |
| SB 04 TN | 4 | 0 | 0.17 | 0.11 | 0.055 | 0.17 |
| SB 00 TDN | 5 | 0 | 0.25 | 0.031 | 0.014 | 0.038 |
| SB 01 TDN | 13 | 0 | 0.20 | 0.081 | 0.022 | 0.049 |
| SB 02 TDN | 11 | 0 | 0.19 | 0.088 | 0.026 | 0.059 |
| SB 04 TDN | 4 | 0 | 0.15 | 0.088 | 0.044 | 0.14 |

| Column | Range | Max | Min | Median | 25% | 75% |
|-----------|--------|--------|---------|--------|--------|--------|
| SB 97 NOx | 0.012 | 0.017 | 0.0050 | 0.0050 | 0.0050 | 0.0050 |
| SB 98 NOx | 0.0040 | 0.0050 | 0.00100 | 0.0050 | 0.0050 | 0.0050 |
| SB 99 NOx | 0.020 | 0.025 | 0.0050 | 0.0050 | 0.0050 | 0.015 |
| SB 00 NOx | 0.051 | 0.056 | 0.0050 | 0.0080 | 0.0050 | 0.037 |
| SB 01 NOx | 0.058 | 0.063 | 0.0050 | 0.0090 | 0.0065 | 0.049 |
| SB 02 NOx | 0.054 | 0.062 | 0.0080 | 0.020 | 0.0100 | 0.033 |
| SB 04 NOx | 0.042 | 0.047 | 0.0050 | 0.024 | 0.014 | 0.036 |
| SB 00 TN | 0.14 | 0.34 | 0.20 | 0.24 | 0.22 | 0.28 |
| SB 01 TN | 0.27 | 0.39 | 0.12 | 0.18 | 0.16 | 0.26 |
| SB 02 TN | 0.24 | 0.35 | 0.11 | 0.16 | 0.12 | 0.23 |
| SB 04 TN | 0.22 | 0.27 | 0.050 | 0.19 | 0.080 | 0.27 |
| SB 00 TDN | 0.080 | 0.29 | 0.21 | 0.26 | 0.23 | 0.27 |
| SB 01 TDN | 0.24 | 0.35 | 0.11 | 0.18 | 0.13 | 0.24 |
| SB 02 TDN | 0.26 | 0.35 | 0.090 | 0.16 | 0.13 | 0.22 |
| SB 04 TDN | 0.21 | 0.26 | 0.050 | 0.15 | 0.085 | 0.22 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|-----------|----------|----------|-----------|-----------|-------|----------------|
| SB 97 NOx | 3.38 | 11.89 | 0.50 | <0.001 | 0.14 | 0.0010 |
| SB 98 NOx | -3.87 | 15.00 | 0.54 | <0.001 | 0.071 | 0.00035 |
| SB 99 NOx | 1.24 | 0.19 | 0.41 | <0.001 | 0.14 | 0.0020 |
| SB 00 NOx | 1.19 | -0.51 | 0.33 | 0.001 | 0.21 | 0.0082 |
| SB 01 NOx | 0.77 | -1.36 | 0.34 | <0.001 | 0.31 | 0.014 |
| SB 02 NOx | 1.39 | 2.11 | 0.18 | 0.399 | 0.26 | 0.0086 |
| SB 04 NOx | 0.34 | 1.30 | 0.23 | 0.533 | 0.100 | 0.0034 |
| SB 00 TN | 1.31 | 2.28 | 0.25 | 0.327 | 1.27 | 0.33 |
| SB 01 TN | 1.08 | 0.017 | 0.28 | 0.007 | 2.81 | 0.70 |
| SB 02 TN | 1.24 | 0.30 | 0.26 | 0.035 | 2.06 | 0.46 |
| SB 04 TN | -0.25 | -4.52 | 0.29 | 0.272 | 0.69 | 0.16 |

| | | | | | | |
|-----------|--------|-------|------|-------|------|------|
| SB 00 TDN | -0.085 | -0.66 | 0.23 | 0.464 | 1.25 | 0.32 |
| SB 01 TDN | 0.81 | -0.56 | 0.18 | 0.321 | 2.61 | 0.60 |
| SB 02 TDN | 1.13 | 0.26 | 0.21 | 0.187 | 2.05 | 0.46 |
| SB 04 TDN | 0.30 | 0.054 | 0.16 | 0.709 | 0.60 | 0.11 |

One Way Analysis of Variance

Data source: SB Water Quality Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: SB in Water Quality Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|-----------|----|---------|--------|--------|--------|
| SB 97 NOx | 24 | 0 | 0.0050 | 0.0050 | 0.0050 |
| SB 98 NOx | 15 | 0 | 0.0050 | 0.0050 | 0.0050 |
| SB 99 NOx | 15 | 0 | 0.0050 | 0.0050 | 0.015 |
| SB 00 NOx | 11 | 0 | 0.0080 | 0.0050 | 0.037 |
| SB 01 NOx | 13 | 0 | 0.0090 | 0.0065 | 0.049 |
| SB 02 NOx | 11 | 0 | 0.020 | 0.0100 | 0.033 |
| SB 04 NOx | 4 | 0 | 0.024 | 0.014 | 0.036 |

H = 42.54 with 6 degrees of freedom. (P = <0.001)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = <0.001)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|------------------------|---------------|------|--------|
| SB 02 NOx vs SB 98 NOx | 47.61 | 4.44 | Yes |
| SB 02 NOx vs SB 97 NOx | 40.79 | 4.15 | Yes |
| SB 02 NOx vs SB 99 NOx | 30.71 | 2.87 | No |
| SB 02 NOx vs SB 00 NOx | 20.05 | 1.74 | No |
| SB 02 NOx vs SB 01 NOx | 10.50 | 0.95 | No |
| SB 02 NOx vs SB 04 NOx | 6.40 | 0.41 | No |
| SB 04 NOx vs SB 98 NOx | 41.21 | 2.71 | No |
| SB 04 NOx vs SB 97 NOx | 34.40 | 2.36 | No |
| SB 04 NOx vs SB 99 NOx | 24.31 | 1.60 | No |
| SB 04 NOx vs SB 00 NOx | 13.65 | 0.87 | No |
| SB 04 NOx vs SB 01 NOx | 4.11 | 0.27 | No |
| SB 01 NOx vs SB 98 NOx | 37.10 | 3.63 | No |
| SB 01 NOx vs SB 97 NOx | 30.29 | 3.26 | No |
| SB 01 NOx vs SB 99 NOx | 20.20 | 1.98 | No |
| SB 01 NOx vs SB 00 NOx | 9.54 | 0.86 | No |
| SB 00 NOx vs SB 98 NOx | 27.56 | 2.57 | No |

| | | | |
|------------------------|-------|------|----|
| SB 00 NOx vs SB 97 NOx | 20.75 | 2.11 | No |
| SB 00 NOx vs SB 99 NOx | 10.66 | 0.99 | No |
| SB 99 NOx vs SB 98 NOx | 16.90 | 1.71 | No |
| SB 99 NOx vs SB 97 NOx | 10.09 | 1.14 | No |
| SB 97 NOx vs SB 98 NOx | 6.81 | 0.77 | No |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

One Way Analysis of Variance

Data source: SB Water Quality Trend Analysis

Normality Test: Failed (P = 0.005)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: SB in Water Quality Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|----------|----|---------|--------|-------|------|
| SB 00 TN | 5 | 0 | 0.24 | 0.22 | 0.28 |
| SB 01 TN | 13 | 0 | 0.18 | 0.16 | 0.26 |
| SB 02 TN | 11 | 0 | 0.16 | 0.12 | 0.23 |
| SB 04 TN | 4 | 0 | 0.19 | 0.080 | 0.27 |

H = 3.90 with 3 degrees of freedom. (P = 0.273)

The differences in the median values among the treatment groups are not great enough to exclude the possibility that the difference is due to random sampling variability; there is not a statistically significant difference (P = 0.273)

One Way Analysis of Variance

Data source: SB Water Quality Trend Analysis

Normality Test: Passed (P = 0.097)

Equal Variance Test: Passed (P = 0.824)

| Group Name | N | Missing | Mean | Std Dev | SEM |
|------------|----|---------|------|---------|-------|
| SB 00 TDN | 5 | 0 | 0.25 | 0.031 | 0.014 |
| SB 01 TDN | 13 | 0 | 0.20 | 0.081 | 0.022 |
| SB 02 TDN | 11 | 0 | 0.19 | 0.088 | 0.026 |
| SB 04 TDN | 4 | 0 | 0.15 | 0.088 | 0.044 |

| Source of Variation | DF | SS | MS | F | P |
|---------------------|----|-------|--------|------|-------|
| Between Groups | 3 | 0.024 | 0.0081 | 1.29 | 0.297 |
| Residual | 29 | 0.18 | 0.0063 | | |
| Total | 32 | 0.21 | | | |

The differences in the mean values among the treatment groups are not great enough to exclude the possibility that the difference is due to random sampling variability; there is not a statistically significant difference (P = 0.297).

Power of performed test with alpha = 0.05: 0.097

The power of the performed test (0.097) is below the desired power of 0.800.
 You should interpret the negative findings cautiously.

Descriptive Statistics:

Data source: SB Eelgrass Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|---------|------|---------|--------|---------|------------|--------------|
| SB 1999 | 12 | 0 | 805.33 | 237.65 | 68.60 | 150.99 |
| SB 2000 | 60 | 0 | 471.17 | 238.09 | 30.74 | 61.50 |
| SB 2001 | 60 | 0 | 466.83 | 247.46 | 31.95 | 63.93 |
| SB 2002 | 60 | 0 | 384.33 | 120.71 | 15.58 | 31.18 |
| SB 2004 | 60 | 0 | 209.83 | 180.35 | 23.28 | 46.59 |

| Column | Range | Max | Min | Median | 25% | 75% |
|---------|--------|---------|--------|--------|--------|--------|
| SB 1999 | 864.00 | 1392.00 | 528.00 | 768.00 | 632.00 | 864.00 |
| SB 2000 | 930.00 | 1070.00 | 140.00 | 420.00 | 300.00 | 540.00 |
| SB 2001 | 950.00 | 970.00 | 20.00 | 405.00 | 285.00 | 685.00 |
| SB 2002 | 470.00 | 660.00 | 190.00 | 370.00 | 285.00 | 455.00 |
| SB 2004 | 760.00 | 760.00 | 0.00 | 190.00 | 50.00 | 290.00 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|---------|----------|----------|-----------|-----------|----------|----------------|
| SB 1999 | 1.46 | 2.54 | 0.24 | 0.064 | 9664.00 | 8403968.00 |
| SB 2000 | 0.99 | 0.36 | 0.15 | 0.001 | 28270.00 | 16664300.00 |
| SB 2001 | 0.27 | -0.85 | 0.12 | 0.027 | 28010.00 | 16688900.00 |
| SB 2002 | 0.49 | -0.53 | 0.083 | 0.372 | 23060.00 | 9722400.00 |
| SB 2004 | 0.79 | 0.22 | 0.12 | 0.026 | 12590.00 | 4560900.00 |

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: SB Eelgrass Trend Analysis

Normality Test: Failed (P = <0.001)

| Group | N | Missing | Median | 25% | 75% |
|---------|----|---------|--------|--------|--------|
| SB 1999 | 12 | 0 | 768.00 | 632.00 | 864.00 |
| SB 2000 | 60 | 0 | 420.00 | 300.00 | 540.00 |
| SB 2001 | 60 | 0 | 405.00 | 285.00 | 685.00 |
| SB 2002 | 60 | 0 | 370.00 | 285.00 | 455.00 |
| SB 2004 | 60 | 0 | 190.00 | 50.00 | 290.00 |

H = 72.06 with 4 degrees of freedom. (P = <0.001)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = <0.001)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|--------------------|---------------|------|--------|
| SB 1999 vs SB 2004 | 159.27 | 6.91 | Yes |
| SB 1999 vs SB 2002 | 97.38 | 4.22 | Yes |

| | | | |
|--------------------|-------|------|-----|
| SB 1999 vs SB 2001 | 79.96 | 3.47 | Yes |
| SB 1999 vs SB 2000 | 78.49 | 3.41 | Yes |
| SB 2000 vs SB 2004 | 80.78 | 6.07 | Yes |
| SB 2000 vs SB 2002 | 18.89 | 1.42 | No |
| SB 2000 vs SB 2001 | 1.47 | 0.11 | No |
| SB 2001 vs SB 2004 | 79.31 | 5.96 | Yes |
| SB 2001 vs SB 2002 | 17.42 | 1.31 | No |
| SB 2002 vs SB 2004 | 61.88 | 4.65 | Yes |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

Descriptive Statistics:

Data source: SB Algae Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|---------|------|---------|-------|---------|------------|--------------|
| SB 2000 | 24 | 0 | 1.67 | 4.82 | 0.98 | 2.03 |
| SB 2001 | 60 | 0 | 63.67 | 32.15 | 4.15 | 8.31 |
| SB 2002 | 60 | 0 | 32.62 | 36.23 | 4.68 | 9.36 |
| SB 2004 | 60 | 1 | 34.25 | 36.94 | 4.81 | 9.63 |

| Column | Range | Max | Min | Median | 25% | 75% |
|---------|--------|--------|-------|--------|-------|--------|
| SB 2000 | 20.00 | 20.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| SB 2001 | 90.00 | 100.00 | 10.00 | 60.00 | 30.00 | 100.00 |
| SB 2002 | 100.00 | 100.00 | 0.00 | 10.00 | 0.00 | 50.00 |
| SB 2004 | 100.00 | 100.00 | 0.00 | 25.00 | 1.00 | 57.50 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|---------|----------|----------|-----------|-----------|---------|----------------|
| SB 2000 | 3.07 | 9.46 | 0.51 | <0.001 | 40.00 | 600.00 |
| SB 2001 | -0.18 | -1.45 | 0.20 | <0.001 | 3820.00 | 304200.00 |
| SB 2002 | 0.72 | -0.88 | 0.25 | <0.001 | 1957.00 | 141257.00 |
| SB 2004 | 0.81 | -0.84 | 0.19 | <0.001 | 2021.00 | 148369.00 |

One Way Analysis of Variance

Data source: SB Algae Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: SB in Algae Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|---------|----|---------|--------|-------|--------|
| SB 2000 | 24 | 0 | 0.00 | 0.00 | 0.00 |
| SB 2001 | 60 | 0 | 60.00 | 30.00 | 100.00 |
| SB 2002 | 60 | 0 | 10.00 | 0.00 | 50.00 |
| SB 2004 | 60 | 1 | 25.00 | 1.00 | 57.50 |

H = 66.25 with 3 degrees of freedom. (P = <0.001)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference ($P = <0.001$)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|--------------------|---------------|------|--------|
| SB 2001 vs SB 2000 | 109.67 | 7.73 | Yes |
| SB 2001 vs SB 2002 | 52.37 | 4.88 | Yes |
| SB 2001 vs SB 2004 | 44.49 | 4.13 | Yes |
| SB 2004 vs SB 2000 | 65.17 | 4.58 | Yes |
| SB 2004 vs SB 2002 | 7.87 | 0.73 | No |
| SB 2002 vs SB 2000 | 57.30 | 4.04 | Yes |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

Appendix 11. The statistical reports generated for all of the analyses conducted in the synthesis of 2004 Long-Term Monitoring Report for Three Mile Harbor.

Descriptive Statistics:

Data source: TMH Water Quality Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|------------------------|------|---------|--------|---------|------------|--------------|
| TMH 97 NO _x | 18 | 0 | 0.0057 | 0.0021 | 0.00050 | 0.0010 |
| TMH 98 NO _x | 13 | 0 | 0.0072 | 0.0059 | 0.0016 | 0.0036 |
| TMH 99 NO _x | 16 | 0 | 0.0081 | 0.0064 | 0.0016 | 0.0034 |
| TMH 00 NO _x | 18 | 0 | 0.021 | 0.030 | 0.0071 | 0.015 |
| TMH 01 NO _x | 16 | 0 | 0.052 | 0.041 | 0.010 | 0.022 |
| TMH 02 NO _x | 17 | 0 | 0.021 | 0.022 | 0.0053 | 0.011 |
| TMH 04 NO _x | 6 | 0 | 0.028 | 0.039 | 0.016 | 0.041 |
| TMH 00 TN | 9 | 0 | 0.28 | 0.048 | 0.016 | 0.037 |
| TMH 01 TN | 16 | 0 | 0.22 | 0.085 | 0.021 | 0.045 |
| TMH 02 TN | 17 | 0 | 0.19 | 0.090 | 0.022 | 0.046 |
| TMH 04 TN | 6 | 0 | 0.18 | 0.11 | 0.044 | 0.11 |
| TMH 00 TDN | 9 | 0 | 0.28 | 0.046 | 0.015 | 0.036 |
| TMH 01 TDN | 16 | 0 | 0.20 | 0.081 | 0.020 | 0.043 |
| TMH 02 TDN | 17 | 0 | 0.19 | 0.082 | 0.020 | 0.042 |
| TMH 04 TDN | 6 | 0 | 0.18 | 0.12 | 0.050 | 0.13 |

| Column | Range | Max | Min | Median | 25% | 75% |
|------------------------|--------|-------|--------|--------|--------|--------|
| TMH 97 NO _x | 0.0070 | 0.012 | 0.0050 | 0.0050 | 0.0050 | 0.0050 |
| TMH 98 NO _x | 0.020 | 0.025 | 0.0050 | 0.0050 | 0.0050 | 0.0050 |
| TMH 99 NO _x | 0.022 | 0.027 | 0.0050 | 0.0050 | 0.0050 | 0.0075 |
| TMH 00 NO _x | 0.12 | 0.13 | 0.0050 | 0.0060 | 0.0050 | 0.032 |
| TMH 01 NO _x | 0.14 | 0.15 | 0.0050 | 0.043 | 0.018 | 0.075 |
| TMH 02 NO _x | 0.081 | 0.086 | 0.0050 | 0.011 | 0.0050 | 0.028 |
| TMH 04 NO _x | 0.10 | 0.11 | 0.0050 | 0.013 | 0.0070 | 0.022 |
| TMH 00 TN | 0.14 | 0.32 | 0.18 | 0.29 | 0.26 | 0.32 |
| TMH 01 TN | 0.27 | 0.39 | 0.12 | 0.19 | 0.17 | 0.26 |
| TMH 02 TN | 0.31 | 0.39 | 0.080 | 0.19 | 0.12 | 0.25 |
| TMH 04 TN | 0.26 | 0.34 | 0.080 | 0.14 | 0.100 | 0.28 |
| TMH 00 TDN | 0.15 | 0.33 | 0.18 | 0.27 | 0.26 | 0.32 |
| TMH 01 TDN | 0.24 | 0.37 | 0.13 | 0.17 | 0.16 | 0.21 |
| TMH 02 TDN | 0.29 | 0.38 | 0.090 | 0.17 | 0.11 | 0.24 |
| TMH 04 TDN | 0.29 | 0.34 | 0.050 | 0.14 | 0.070 | 0.31 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|------------------------|----------|----------|-----------|-----------|-------|----------------|
| TMH 97 NO _x | 2.74 | 6.28 | 0.52 | <0.001 | 0.10 | 0.00067 |
| TMH 98 NO _x | 2.79 | 7.77 | 0.49 | <0.001 | 0.094 | 0.0011 |
| TMH 99 NO _x | 2.34 | 5.08 | 0.37 | <0.001 | 0.13 | 0.0016 |
| TMH 00 NO _x | 2.73 | 8.64 | 0.31 | <0.001 | 0.38 | 0.024 |
| TMH 01 NO _x | 1.00 | 0.69 | 0.15 | 0.437 | 0.83 | 0.068 |
| TMH 02 NO _x | 1.97 | 4.03 | 0.27 | 0.002 | 0.35 | 0.015 |
| TMH 04 NO _x | 2.33 | 5.52 | 0.39 | 0.005 | 0.17 | 0.012 |
| TMH 00 TN | -1.32 | 1.24 | 0.20 | 0.338 | 2.52 | 0.72 |
| TMH 01 TN | 1.08 | -0.14 | 0.25 | 0.010 | 3.51 | 0.88 |
| TMH 02 TN | 0.67 | -0.25 | 0.18 | 0.176 | 3.31 | 0.77 |
| TMH 04 TN | 0.80 | -1.34 | 0.27 | 0.203 | 1.07 | 0.25 |
| TMH 00 TDN | -1.14 | 1.81 | 0.23 | 0.182 | 2.51 | 0.72 |
| TMH 01 TDN | 1.52 | 0.81 | 0.30 | <0.001 | 3.25 | 0.76 |
| TMH 02 TDN | 0.87 | 0.25 | 0.16 | 0.266 | 3.15 | 0.69 |
| TMH 04 TDN | 0.62 | -1.71 | 0.28 | 0.152 | 1.05 | 0.26 |

One Way Analysis of Variance

Data source: TMH Water Quality Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: TMH in Water Quality Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|------------------------|----|---------|--------|--------|--------|
| TMH 97 NO _x | 18 | 0 | 0.0050 | 0.0050 | 0.0050 |
| TMH 98 NO _x | 13 | 0 | 0.0050 | 0.0050 | 0.0050 |
| TMH 99 NO _x | 16 | 0 | 0.0050 | 0.0050 | 0.0075 |
| TMH 00 NO _x | 18 | 0 | 0.0060 | 0.0050 | 0.032 |
| TMH 01 NO _x | 16 | 0 | 0.043 | 0.018 | 0.075 |
| TMH 02 NO _x | 17 | 0 | 0.011 | 0.0050 | 0.028 |
| TMH 04 NO _x | 6 | 0 | 0.013 | 0.0070 | 0.022 |

H = 40.09 with 6 degrees of freedom. (P = <0.001)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = <0.001)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|--|---------------|------|--------|
| TMH 01 NO _x vs TMH 97 NO _x | 50.69 | 4.89 | Yes |
| TMH 01 NO _x vs TMH 98 NO _x | 47.44 | 4.21 | Yes |
| TMH 01 NO _x vs TMH 99 NO _x | 41.91 | 3.93 | Yes |
| TMH 01 NO _x vs TMH 00 NO _x | 29.08 | 2.81 | No |
| TMH 01 NO _x vs TMH 02 NO _x | 20.47 | 1.95 | No |
| TMH 01 NO _x vs TMH 04 NO _x | 15.77 | 1.09 | No |
| TMH 04 NO _x vs TMH 97 NO _x | 34.92 | 2.46 | No |
| TMH 04 NO _x vs TMH 98 NO _x | 31.67 | 2.13 | No |
| TMH 04 NO _x vs TMH 99 NO _x | 26.14 | 1.81 | No |
| TMH 04 NO _x vs TMH 00 NO _x | 13.31 | 0.94 | No |
| TMH 04 NO _x vs TMH 02 NO _x | 4.70 | 0.33 | No |
| TMH 02 NO _x vs TMH 97 NO _x | 30.22 | 2.96 | No |
| TMH 02 NO _x vs TMH 98 NO _x | 26.97 | 2.43 | No |
| TMH 02 NO _x vs TMH 99 NO _x | 21.44 | 2.04 | No |
| TMH 02 NO _x vs TMH 00 NO _x | 8.61 | 0.84 | No |
| TMH 00 NO _x vs TMH 97 NO _x | 21.61 | 2.15 | No |
| TMH 00 NO _x vs TMH 98 NO _x | 18.36 | 1.67 | No |
| TMH 00 NO _x vs TMH 99 NO _x | 12.83 | 1.24 | No |
| TMH 99 NO _x vs TMH 97 NO _x | 8.78 | 0.85 | No |
| TMH 99 NO _x vs TMH 98 NO _x | 5.53 | 0.49 | No |

TMH 98 NO_x vs TMH 97 NO_x 3.25 0.30 No

Note: The multiple comparisons on ranks do not include an adjustment for ties.

One Way Analysis of Variance

Data source: TMH Water Quality Trend Analysis

Normality Test: Passed (P > 0.200)

Equal Variance Test: Passed (P = 0.360)

| Group Name | N | Missing | Mean | Std Dev | SEM |
|------------|----|---------|------|---------|-------|
| TMH 00 TN | 9 | 0 | 0.28 | 0.048 | 0.016 |
| TMH 01 TN | 16 | 0 | 0.22 | 0.085 | 0.021 |
| TMH 02 TN | 17 | 0 | 0.19 | 0.090 | 0.022 |
| TMH 04 TN | 6 | 0 | 0.18 | 0.11 | 0.044 |

| Source of Variation | DF | SS | MS | F | P |
|---------------------|----|-------|--------|------|-------|
| Between Groups | 3 | 0.053 | 0.018 | 2.48 | 0.074 |
| Residual | 44 | 0.32 | 0.0072 | | |
| Total | 47 | 0.37 | | | |

The differences in the mean values among the treatment groups are not great enough to exclude the possibility that the difference is due to random sampling variability; there is not a statistically significant difference (P = 0.074).

Power of performed test with alpha = 0.05: 0.354

The power of the performed test (0.354) is below the desired power of 0.800.

You should interpret the negative findings cautiously.

One Way Analysis of Variance

Data source: TMH Water Quality Trend Analysis

Normality Test: Failed (P = 0.002)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: TMH in Water Quality Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|------------|----|---------|--------|-------|------|
| TMH 00 TDN | 9 | 0 | 0.27 | 0.26 | 0.32 |
| TMH 01 TDN | 16 | 0 | 0.17 | 0.16 | 0.21 |
| TMH 02 TDN | 17 | 0 | 0.17 | 0.11 | 0.24 |
| TMH 04 TDN | 6 | 0 | 0.14 | 0.070 | 0.31 |

H = 9.24 with 3 degrees of freedom. (P = 0.026)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = 0.026)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|--------------------------|---------------|------|--------|
| TMH 00 TDN vs TMH 04 TDN | 18.22 | 2.47 | No |
| TMH 00 TDN vs TMH 02 TDN | 16.03 | 2.78 | No |
| TMH 00 TDN vs TMH 01 TDN | 12.31 | 2.11 | No |
| TMH 01 TDN vs TMH 04 TDN | 5.92 | 0.88 | No |
| TMH 01 TDN vs TMH 02 TDN | 3.72 | 0.76 | No |
| TMH 02 TDN vs TMH 04 TDN | 2.20 | 0.33 | No |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

Descriptive Statistics:

Data source: TMH Eelgrass Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|----------|------|---------|--------|---------|------------|--------------|
| TMH 1999 | 12 | 0 | 361.33 | 169.12 | 48.82 | 107.45 |
| TMH 2000 | 60 | 0 | 192.83 | 129.80 | 16.76 | 33.53 |
| TMH 2001 | 60 | 0 | 208.83 | 99.12 | 12.80 | 25.60 |
| TMH 2002 | 60 | 0 | 135.38 | 74.00 | 9.55 | 19.12 |
| TMH 2004 | 60 | 0 | 29.00 | 47.25 | 6.10 | 12.21 |

| Column | Range | Max | Min | Median | 25% | 75% |
|----------|--------|--------|-------|--------|--------|--------|
| TMH 1999 | 480.00 | 576.00 | 96.00 | 376.00 | 208.00 | 504.00 |
| TMH 2000 | 560.00 | 600.00 | 40.00 | 165.00 | 100.00 | 255.00 |
| TMH 2001 | 450.00 | 470.00 | 20.00 | 205.00 | 140.00 | 265.00 |
| TMH 2002 | 260.00 | 260.00 | 0.00 | 150.00 | 110.00 | 190.00 |
| TMH 2004 | 210.00 | 210.00 | 0.00 | 0.00 | 0.00 | 40.00 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|----------|----------|----------|-----------|-----------|----------|----------------|
| TMH 1999 | -0.29 | -1.23 | 0.14 | 0.616 | 4336.00 | 1881344.00 |
| TMH 2000 | 1.31 | 1.44 | 0.16 | <0.001 | 11570.00 | 3225100.00 |
| TMH 2001 | 0.29 | -0.13 | 0.069 | 0.635 | 12530.00 | 3196300.00 |
| TMH 2002 | -0.66 | -0.36 | 0.15 | 0.001 | 8123.00 | 1422809.00 |
| TMH 2004 | 1.96 | 3.71 | 0.31 | <0.001 | 1740.00 | 182200.00 |

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: TMH Eelgrass Trend Analysis

Normality Test: Failed (P = <0.001)

| Group | N | Missing | Median | 25% | 75% |
|----------|----|---------|--------|--------|--------|
| TMH 1999 | 12 | 0 | 376.00 | 208.00 | 504.00 |
| TMH 2000 | 60 | 0 | 165.00 | 100.00 | 255.00 |
| TMH 2001 | 60 | 0 | 205.00 | 140.00 | 265.00 |
| TMH 2002 | 60 | 0 | 150.00 | 110.00 | 190.00 |
| TMH 2004 | 60 | 0 | 0.00 | 0.00 | 40.00 |

H = 115.57 with 4 degrees of freedom. (P = <0.001)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = <0.001)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|----------------------|---------------|------|-------------|
| TMH 1999 vs TMH 2004 | 163.83 | 7.11 | Yes |
| TMH 1999 vs TMH 2002 | 83.64 | 3.63 | Yes |
| TMH 1999 vs TMH 2000 | 60.57 | 2.63 | No |
| TMH 1999 vs TMH 2001 | 41.07 | 1.78 | Do Not Test |
| TMH 2001 vs TMH 2004 | 122.76 | 9.22 | Yes |
| TMH 2001 vs TMH 2002 | 42.57 | 3.20 | Yes |
| TMH 2001 vs TMH 2000 | 19.50 | 1.47 | Do Not Test |
| TMH 2000 vs TMH 2004 | 103.26 | 7.76 | Yes |
| TMH 2000 vs TMH 2002 | 23.07 | 1.73 | No |
| TMH 2002 vs TMH 2004 | 80.19 | 6.03 | Yes |

Note: The multiple comparisons on ranks do not include an adjustment for ties.

Descriptive Statistics:

Data source: TMH Algae Trend Analysis

| Column | Size | Missing | Mean | Std Dev | Std. Error | C.I. of Mean |
|----------|------|---------|-------|---------|------------|--------------|
| TMH 2000 | 24 | 0 | 47.08 | 42.37 | 8.65 | 17.89 |
| TMH 2001 | 60 | 0 | 38.37 | 35.15 | 4.54 | 9.08 |
| TMH 2002 | 60 | 0 | 22.83 | 29.12 | 3.76 | 7.52 |
| TMH 2004 | 60 | 2 | 14.72 | 30.11 | 3.95 | 7.92 |

| Column | Range | Max | Min | Median | 25% | 75% |
|----------|--------|--------|------|--------|------|-------|
| TMH 2000 | 100.00 | 100.00 | 0.00 | 45.00 | 0.00 | 90.00 |
| TMH 2001 | 100.00 | 100.00 | 0.00 | 30.00 | 0.00 | 75.00 |
| TMH 2002 | 100.00 | 100.00 | 0.00 | 5.00 | 0.00 | 50.00 |
| TMH 2004 | 100.00 | 100.00 | 0.00 | 0.00 | 0.00 | 10.00 |

| Column | Skewness | Kurtosis | K-S Dist. | K-S Prob. | Sum | Sum of Squares |
|----------|----------|----------|-----------|-----------|---------|----------------|
| TMH 2000 | 0.072 | -1.81 | 0.20 | 0.014 | 1130.00 | 94500.00 |
| TMH 2001 | 0.41 | -1.16 | 0.17 | <0.001 | 2302.00 | 161202.00 |
| TMH 2002 | 1.00 | -0.38 | 0.29 | <0.001 | 1370.00 | 81300.00 |
| TMH 2004 | 2.11 | 3.06 | 0.36 | <0.001 | 854.00 | 64238.00 |

One Way Analysis of Variance

Data source: TMH Algae Trend Analysis

Normality Test: Failed (P = <0.001)

Test execution ended by user request, ANOVA on Ranks begun

Kruskal-Wallis One Way Analysis of Variance on Ranks

Data source: TMH in Algae Trend Analysis

| Group | N | Missing | Median | 25% | 75% |
|----------|----|---------|--------|------|-------|
| TMH 2000 | 24 | 0 | 45.00 | 0.00 | 90.00 |
| TMH 2001 | 60 | 0 | 30.00 | 0.00 | 75.00 |
| TMH 2002 | 60 | 0 | 5.00 | 0.00 | 50.00 |
| TMH 2004 | 60 | 2 | 0.00 | 0.00 | 10.00 |

H = 20.03 with 3 degrees of freedom. (P = <0.001)

The differences in the median values among the treatment groups are greater than would be expected by chance; there is a statistically significant difference (P = <0.001)

To isolate the group or groups that differ from the others use a multiple comparison procedure.

All Pairwise Multiple Comparison Procedures (Dunn's Method) :

| Comparison | Diff of Ranks | Q | P<0.05 |
|----------------------|---------------|------|--------|
| TMH 2000 vs TMH 2004 | 46.50 | 3.28 | Yes |
| TMH 2000 vs TMH 2002 | 28.91 | 2.05 | No |
| TMH 2000 vs TMH 2001 | 6.24 | 0.44 | No |
| TMH 2001 vs TMH 2004 | 40.26 | 3.74 | Yes |
| TMH 2001 vs TMH 2002 | 22.67 | 2.12 | No |
| TMH 2002 vs TMH 2004 | 17.59 | 1.63 | No |

Note: The multiple comparisons on ranks do not include an adjustment for ties.